BOUNDARY MINIMAL MODELS AND THE ROGERS-RAMANUJAN IDENTITIES

DIEGO SALAZAR

ABSTRACT. We determine when the irreducible modules $L(c_{p,q}, h_{m,n})$ over the simple Virasoro vertex algebras $\operatorname{Vir}_{p,q}$, where $p, q \geq 2$ are relatively prime with 0 < m < p and 0 < n < q, are classically free. It turns out this only happens with the boundary minimal models, i.e., with the irreducible modules over $\operatorname{Vir}_{2,2s+1}$ for $s \in \mathbb{Z}_+$. We thus obtain a complete description of the classical limits of these modules in terms of the jet algebra of the corresponding Zhu C_2 -algebra. Gordon's generalization of the Rogers-Ramanujan identities is used in the proof, and our results in turn provide a natural interpretation of these identities.

Keywords. Vertex algebras, boundary minimal models, Rogers-Ramanujan identities, quantum algebra, combinatorics.

0. INTRODUCTION

In [Li05], Li introduced a decreasing filtration $(F_pV)_{p\in\mathbb{Z}}$ on an arbitrary vertex algebra V. The associated graded space $\operatorname{gr}_F(V)$ with respect to this decreasing filtration carries the structure of a vertex Poisson algebra. Li also introduced a decreasing filtration $(F_pM)_{p\in\mathbb{Z}}$ for modules M over a vertex algebra V and showed that the associated graded space $\operatorname{gr}_F(M)$ is a module over the vertex Poisson algebra $\operatorname{gr}_F(V)$.

Given a commutative algebra R, the jet algebra JR is constructed as the smallest differential algebra containing R. It was proved by Arakawa in [Ara12, Proposition 2.5.1] that there exists a natural surjection

$$JR_V \twoheadrightarrow \operatorname{gr}_F(V)$$

of vertex Poisson algebras, where R_V is the Zhu C_2 -algebra of V as defined in [Zhu96]. When this is an isomorphism, we say V is classically free. For modules M over the vertex algebra V, we also have a natural surjection

$$JR_V \otimes_{R_V} R_M \twoheadrightarrow \operatorname{gr}_F(M),$$

and we say M is classically free if this is an isomorphism. One may in turn regard the classically free vertex algebra V or V-module M, as a quantization of JR_V or $JR_V \otimes_{R_V} R_M$, respectively.

The question of classical freedom of the simple Virasoro vertex algebra $\operatorname{Vir}_{p,q}$ was studied in [VEH21], and the answer is very simple. If $q > p \geq 2$ are relatively prime integers, then $\operatorname{Vir}_{p,q}$ is classically free if and only if p = 2. The main objective of this article is to solve this problem for all irreducible modules over the vertex algebras $\operatorname{Vir}_{p,q}$, namely $L(c_{p,q}, h_{m,n})$ for integers m, n such that 0 < m < p and 0 < n < q.

It turns out this problem is deeply connected with the Rogers-Ramanujan q-series identities and Gordon's generalization [Gor61], namely

$$\sum_{k=(k_1,\dots,k_{s-1})\in\mathbb{N}^{s-1}}\frac{q^{\frac{1}{2}kG^{(s)}k^T+kB^{(s)}_{s-i}}}{(q)_{k_1}\dots(q)_{k_{s-1}}} = \prod_{\substack{n\neq 0,\pm i \mod 2s+1}}^{\infty} (1-q^n)^{-1},$$
(0.1)

for $s \in \mathbb{Z}_+$, $i \in \{1, \ldots, s\}$, where

$$G^{(s)} = (2\min\{i, j\})_{i,j=1}^{s-1},$$

$$B_j^{(s)} = (0, 0, \dots, 0, 1, 2, \dots, j)^T \text{ for } j = 0, 1, \dots, s-1,$$

Date: May 12, 2024.

and T denotes the transpose matrix.

The identity (0.1) has an interpretation in terms of integer partitions, as explained in [And98, §8]. We represent a partition λ either as a finite nonincreasing sequence of positive integers $[\lambda_1, \ldots, \lambda_s]$ or else as an infinite sequence $(f_n)_{n \in \mathbb{Z}_+} = [1^{f_1}, 2^{f_2}, \ldots]$ of nonnegative integers in which only finitely many of the f_n are nonzero corresponding, the term f_n being, of course, the number of times n appears among the λ_i . We have a natural partial order \leq on partitions given by $(g_n)_{n \in \mathbb{Z}_+} \leq (f_n)_{n \in \mathbb{Z}_+}$ if $g_n \leq f_n$ for $n \in \mathbb{Z}_+$. A set P of partitions is said to be a partition ideal if whenever $(f_n)_{n \in \mathbb{Z}_+} \in P$ and $(g_n)_{n \in \mathbb{Z}_+} \leq (f_n)_{n \in \mathbb{Z}_+} \in P$. We denote by p(P, n) the number of partitions of n that belong to P. We say two partition ideals P_1 and P_2 are equivalent, denoted by $P_1 \sim P_2$, if $p(P_1, n) = p(P_2, n)$ for $n \in \mathbb{N}$.

Let $s \in \mathbb{Z}_+$, let $i \in \{1, \ldots, s\}$. Gordon's generalization of the Rogers-Ramanujan identities asserts

$$A^{s,i} \sim P^{s,i}$$

where the partition ideals $A^{s,i}$ and $P^{s,i}$ are defined by:

 $A^{s,i} = \{(f_n)_{n \in \mathbb{Z}_+} \text{ is a partition } | \text{ for } n \in \mathbb{Z}_+, f_n > 0 \text{ implies } n \neq 0, \pm i \mod 2s + 1\},\$ $R^{s,i} = \{\lambda \text{ is a partition } | \lambda = [\lambda_1, \dots, \lambda_s] \text{ satisfies } \lambda_1 - \lambda_s \leq 1 \text{ or } \lambda = [1^i]\},\$ $P^{s,i} = \text{the set of partitions that do not contain any partition in } R^{s,i}.$

As explained in [And98, §8], $A^{s,i}$ is a partition ideal of order 1 while $P^{s,i}$ is a partition ideal of order 2. It is an open problem to characterize the equivalence classes of partitions ideals. For partition ideals of order 1, this problem is completely solved (see [And98, Theorem 8.4]).

In the context of representations of infinite dimensional Lie algebras and vertex algebras, the left-hand side of (0.1) is precisely the graded dimension, or character, of the irreducible Vir-module $M = L(c_{2,2s+1}, h_{1,i})$. Li's filtration is compatible with the grading, and so $\operatorname{gr}_F(M)$ acquires a second grading and hence a refined character, a series in two variables (t,q). Our main theorems concern a modified version of this refined character and the intimate relation between the structure of $\operatorname{gr}_F(M)$ and the partition ideals $P^{s,i}$.

Theorem 0.1. The refined character of $L(c_{2,2s+1}, h_{1,i})$ is given by

$$\operatorname{ch}_{L(c_{2,2s+1},h_{1,i})}(t,q) = q^{h_{1,i}} \left(\sum_{k=(k_1,\dots,k_{s-1})\in\mathbb{N}^{s-1}} t^{kB_{s-1}^{(s)}} \frac{q^{\frac{1}{2}kG^{(s)}k^T + kB_{s-i}^{(s)}}}{(q)_{k_1}\dots(q)_{k_{s-1}}} \right).$$

Theorem 0.2. The set

$$L_{-\lambda_1}L_{-\lambda_2}\dots L_{-\lambda_m}|c_{2,2s+1},h_{1,i}\rangle \mid \lambda = [\lambda_1,\dots,\lambda_m] \in P^{s,i}\}$$

is a vector space basis of $L(c_{2,2s+1}, h_{1,i})$.

{

Theorem 0.3. The boundary minimal models $L(c_{2,2s+1}, h_{1,i})$ are classically free as modules over the simple Virasoro vertex algebra $\operatorname{Vir}_{2,2s+1}$ for $s \in \mathbb{Z}_+$ and $i \in \{1, \ldots, s\}$. If p, q > 2 then no irreducible $\operatorname{Vir}_{p,q}$ -module is classically free.

As a corollary, we obtain an explicit expression for the modules $\operatorname{gr}_F(L(c_{2,2s+1}, h_{1,i}))$ over the vertex Poisson algebra $\operatorname{gr}_F(\operatorname{Vir}_{2,2s+1})$.

This article is organized as follows. In §1, we review Verma modules and set notations related to partitions. In §2, we review certain filtrations of algebras and modules, the main example being the PBW filtration of a Lie algebra. In §3, we review modules over Lie algebras graded by a free abelian group of finite rank, the main example being Verma module M(c, h) and the irreducible highest weight representation L(c, h). In §5, we introduce an explicit formula related to the normal product of a finite number of fields. In §6, we recall the irreducible modules over the simple Virasoro vertex algebras. In §7, we summarize the theory of filtrations of vertex algebras and their modules, with emphasis on explicit examples related to modules over the Virasoro Lie algebra. In §8, we recall the Zhu's algebra and the notion of classically freeness of vertex algebras and their modules. In §9, we prove our desired theorems. In the appendices, we recall notions related to Poisson algebras, jet algebras and vertex Poisson algebras.

I would like to thank Reimundo Heluani and Jethro Van Ekeren for some valuable suggestions. The author is partially supported by FAPERJ grant 201.445/2021.

1. Preliminaries and notation

All vector spaces and all algebras are over \mathbb{C} , the field of complex numbers, unless otherwise stated. The set of natural numbers $\{0, 1, ...\}$ is denoted by \mathbb{N} , the set of integers is denoted by \mathbb{Z} , and the set of positive integers $\{1, 2, ...\}$ is denoted by \mathbb{Z}_+ .

First, we review the theory of representations of the Virasoro Lie algebra following [KRR13]. The *Virasoro Lie algebra* is a Lie algebra given by

$$\operatorname{Vir} = \bigoplus_{n \in \mathbb{Z}} \mathbb{C}L_n \oplus \mathbb{C}C.$$

These elements satisfy the following commutation relations:

$$[L_m, L_n] = (m - n)L_{m+n} + \delta_{m, -n} \frac{m^3 - m}{12}C \quad \text{for } m, n \in \mathbb{Z},$$

[Vir, C] = 0.

Let $c, h \in \mathbb{C}$. We set $\operatorname{Vir}^{\geq} = \bigoplus_{n \in \mathbb{N}} \mathbb{C}L_n \oplus \mathbb{C}C$. We make the subalgebra $\operatorname{Vir}^{\geq}$ of Vir act on \mathbb{C} as follows:

$$L_n 1 = 0$$
 for $n \in \mathbb{Z}_+$, $L_0 1 = h$ and $C 1 = c$.

The Verma representation of Vir with highest weight (c, h) is defined as

$$M(c,h) = \operatorname{Ind}_{\operatorname{Vir}^{\geq}}^{\operatorname{Vir}}(\mathbb{C}) = U(\operatorname{Vir}) \otimes_{U(\operatorname{Vir}^{\geq})} \mathbb{C},$$

where Vir acts by left multiplication. We take $|c, h\rangle = 1 \otimes 1$ as highest weight vector.

A composition (of $n \in \mathbb{N}$) is a sequence $\lambda = [\lambda_1, \ldots, \lambda_m]$ such that $\lambda_i \in \mathbb{Z}_+$ for $i = 1, \ldots, m$ (and $\lambda_1 + \cdots + \lambda_m = n$). We also consider the *empty composition* \emptyset , which is the unique composition of 0. Given a composition $\lambda = [\lambda_1, \ldots, \lambda_m]$ and a permutation $\sigma \in S_m$, we define the composition

$$\lambda \sigma = [\lambda_{\sigma(1)}, \ldots, \lambda_{\sigma(m)}].$$

For a composition $\lambda = [\lambda_1, \ldots, \lambda_m]$, we define

$$L_{\lambda} = L_{-\lambda_1} \dots L_{-\lambda_m} \in U(\text{Vir}),$$

the length of λ as

$$\operatorname{len}(\lambda) = m,$$

the weight of λ as

$$\Delta(\lambda) = \lambda_1 + \dots + \lambda_m,$$

and

$$p_{\lambda} = L_{-\lambda_1} \dots L_{-\lambda_m} \in \mathbb{C}[L_{-1}, L_{-2}, \dots].$$

For a composition λ with exactly *m* elements greater than 1 and exactly *n* elements equal to 1, we define λ' as the composition obtained from λ by removing all 1s. We define the *conformal* length of λ as

$$\operatorname{clen}(\lambda) = 2m + n$$

and

$$u_{\lambda} = p_{\lambda'} L_{-1}^n \in \bigoplus_{k \in \mathbb{N}} \mathbb{C}[L_{-2}, L_{-3}, \dots] L_{-1}^k.$$

A partition (of $n \in \mathbb{N}$) is a composition $\lambda = [\lambda_1, \ldots, \lambda_m]$ (of n) such that $\lambda_1 \geq \cdots \geq \lambda_m$. The notation $\lambda \vdash$ means λ is a partition, and for $n \in \mathbb{N}$, the notation $\lambda \vdash n$ means λ is a partition of n. This notation also tacitly assumes that λ is a partition and not just a composition.

Given a composition λ of length m, there is a unique partition $prt(\lambda)$ obtained by reordering λ , i.e., by considering all $\lambda \sigma$ for $\sigma \in S_m$.

By the PBW theorem, the set

$$\{L_{\lambda}|c,h\rangle \mid \lambda \vdash\}$$

is a vector space basis of M(c, h). The representation M(c, h) has a unique maximal proper subrepresentation J(c, h), and the quotient

$$L(c,h) = M(c,h)/J(c,h)$$

is the *irreducible highest weight representation* of Vir with highest weight (c, h).

2. Modules over algebras with a filtration

Let A be an associative (not necessarily commutative) algebra with unit 1 and filtration $(A^p)_{p\in\mathbb{Z}}$ such that:

(i) $A^p = 0$ for p < 0; (ii) $1 \in A^0$; (iii) $A^0 \subseteq A^1 \subseteq \dots$; (iv) $A = \bigcup_{p \in \mathbb{N}} A^p$; (v) $A^p A^q \subseteq A^{p+q}$ for $p, q \in \mathbb{Z}$.

Let

$$\operatorname{gr}(A) = \bigoplus_{p \in \mathbb{N}} A^p / A^{p-1}$$

be the associated graded vector space. The vector space gr(A) is an associative algebra with unit and multiplication given as follows. For $p, q \in \mathbb{N}$, $a \in A^p$ and $b \in A^q$, we set

$$\gamma^p(a)\gamma^q(b) = \gamma^{p+q}(ab)$$

where $\gamma^p : A^p \to \operatorname{gr}(A)$ is the *principal symbol map*, which is the composition of the natural maps $A^p \to A^p/A^{p-1}$ and $A^p/A^{p-1} \to \operatorname{gr}(A)$. The unit of $\operatorname{gr}(A)$ is $\gamma^0(1)$.

Let ∂ be a derivation of A respecting the filtration $(A^p)_{p \in \mathbb{Z}}$, i.e., it satisfies:

- (i) $\partial(ab) = \partial(a)b + a\partial(b)$ for $a, b \in A$;
- (ii) $\partial(A^p) \subseteq A^p$ for $p \in \mathbb{Z}$.

We can define

$$\begin{aligned} \partial : \operatorname{gr}(A) &\to \operatorname{gr}(A), \\ \partial(\gamma^p(a)) &= \gamma^p(\partial(a)) \quad \text{for } p \in \mathbb{Z} \text{ and } a \in A^p, \end{aligned}$$

and it is a derivation of gr(A).

Example 2.1 (PBW filtration of U(\mathfrak{g})). Let \mathfrak{g} be a Lie algebra. The PBW filtration of $U(\mathfrak{g})$, the universal enveloping algebra of \mathfrak{g} , is given by

$$U(\mathfrak{g})^p = \operatorname{span}\{x_1 x_2 \dots x_s \mid s \le p, x_1, \dots, x_s \in \mathfrak{g}\} \quad \text{for } p \in \mathbb{Z}.$$

This filtration clearly satisfies axioms (i)–(v) above, and $\operatorname{gr}(U(\mathfrak{g}))$ is naturally isomorphic to $S(\mathfrak{g})$, the symmetric algebra of \mathfrak{g} , which is a polynomial algebra (see [Dix96, §2] for details). Furthermore, if ∂ is a derivation of \mathfrak{g} as a Lie algebra, then we can extend ∂ to a derivation of $U(\mathfrak{g})$, and it respects the PBW filtration. Thus, it defines a derivation $\partial : \operatorname{gr}(U(\mathfrak{g})) \to \operatorname{gr}(U(\mathfrak{g}))$.

Let M be an A-module with filtration $(M^p)_{p\in\mathbb{Z}}$ such that:

(i)
$$M^p = 0$$
 for $p < 0$;
(ii) $M^0 \subseteq M^1 \subseteq \dots$;
(iii) $M = \bigcup_{p \in \mathbb{N}} M^p$;
(iv) $A^p M^q \subseteq M^{p+q}$ for $p, q \in \mathbb{Z}$.
Let

$$\operatorname{gr}(M) = \bigoplus_{p \in \mathbb{N}} M^p / M^{p-1}$$

be the associated graded vector space.

Then $\operatorname{gr}(M)$ is a $\operatorname{gr}(A)$ -module with operations given as follows. For $p, q \in \mathbb{N}$, $a \in A^p$ and $u \in M^q$, we set

$$\gamma^p(a)\gamma^q_M(u) = \gamma^{p+q}_M(au),$$

where $\gamma_M^p: M^p \to \operatorname{gr}(M)$ is the principal symbol map, which is the composition of the natural maps $M^p \to M^p/M^{p-1}$ and $M^p/M^{p-1} \hookrightarrow \operatorname{gr}(M)$.

The category of modules over A with the given filtration $(A^p)_{p\in\mathbb{Z}}$ is given by modules M with filtration $(M^p)_{p\in\mathbb{Z}}$ satisfying conditions (i)–(iv) above. A homomorphism $f: M \to N$ must satisfy $f(M^p) \subseteq N^p$ for $p \in \mathbb{Z}$. This category is denoted by A-Mod (the filtration being suppressed from the notation).

If $f: M \to N$ is a homomorphism of A-modules, then

$$\operatorname{gr}(f) : \operatorname{gr}(M) \to \operatorname{gr}(N),$$

 $\operatorname{gr}(f)(\gamma_M^p(u)) = \gamma_N^p(f(u)) \text{ for } p \in \mathbb{N} \text{ and } u \in M^p$

defines a homomorphism of gr(A)-modules. Therefore, we obtain a functor

$$\operatorname{gr} : A\operatorname{-Mod} \to \operatorname{gr}(A)\operatorname{-Mod}.$$

From now on, some subscripts or superscripts will be omitted, so for example γ_M^p simplifies to γ^p .

3. Modules over Q-graded Lie Algebras

In this section, we follow [IK11, §2]. Let Γ be an abelian group. For a Γ -graded vector space $V = \bigoplus_{\alpha \in \Gamma} V^{\alpha}$, we set $\operatorname{supp}(V) = \{ \alpha \in \Gamma \mid V^{\alpha} \neq 0 \}.$

A Γ -graded Lie algebra is a Lie algebra $\mathfrak{g} = \bigoplus_{\alpha \in \Gamma} \mathfrak{g}^{\alpha}$ such that

$$[\mathfrak{g}^{\alpha},\mathfrak{g}^{\beta}]\subseteq\mathfrak{g}^{\alpha+\beta}\quad\text{for }\alpha,\beta\in\Gamma$$

Let Q be a free abelian group of finite rank r, and let \mathfrak{g} be a Lie algebra with a commutative subalgebra \mathfrak{h} . We say a pair $(\mathfrak{g}, \mathfrak{h})$ is a Q-graded Lie algebra if it satisfies the following:

(i) $\mathfrak{g} = \bigoplus_{\alpha \in Q} \mathfrak{g}^{\alpha}$ is Q-graded, $\mathfrak{h} = \mathfrak{g}^{0}$, and $\operatorname{supp}(\mathfrak{g})$ generates Q;

(ii) We have a homomorphism $\pi_Q: Q \to \mathfrak{h}^*, \alpha \mapsto \lambda_\alpha$ such that

$$[h, x] = \lambda_{\alpha}(h)x \text{ for } h \in \mathfrak{h} \text{ and } x \in \mathfrak{g}^{\alpha};$$

(iii) For $\alpha \in Q$, dim $(\mathfrak{g}^{\alpha}) < \infty$;

(iv) There exists a basis $(\alpha_i)_{i=1}^r$ of Q such that for $\alpha \in \operatorname{supp}(\mathfrak{g})$,

$$\alpha \in \sum_{i=1}^{r} \mathbb{Z}_{\geq 0} \alpha_i \text{ or } \alpha \in \sum_{i=1}^{r} \mathbb{Z}_{\leq 0} \alpha_i$$

The condition (iv) implies that a Q-graded Lie algebra admits a triangular decomposition

$$\mathfrak{g} = \mathfrak{g}^- \oplus \mathfrak{h} \oplus \mathfrak{g}^+,$$

where $Q^+ = \sum_{i=1}^r \mathbb{Z}_{\geq 0} \alpha_i$ and

$$\mathfrak{g}^{\pm} = igoplus_{\pm lpha \in Q^+ \setminus \{0\}} \mathfrak{g}^{lpha}$$

For later use, we set $\mathfrak{g}^{\geq} = \mathfrak{h} \oplus \mathfrak{g}^+$ and $\mathfrak{g}^{\leq} = \mathfrak{g}^- \oplus \mathfrak{h}$.

Example 3.1. We set $Q = \mathbb{Z}$, $\mathfrak{h} = \mathbb{C}L_0 \oplus \mathbb{C}C$, $\operatorname{Vir}^n = \mathbb{C}L_n$ for $n \in \mathbb{Z} \setminus \{0\}$ and

$$\pi_Q: Q \to \mathfrak{h}^*, \pi_Q(n) = (L_0 \mapsto -n, C \mapsto 0).$$

Then (Vir, \mathfrak{h}) is readily seen to be a \mathbb{Z} -graded Lie algebra. We take 1 as the basis of Q, so $Q^+ = \mathbb{N}$. We can verify that $\mathrm{ad}(L_{-1})$ is a derivation of Vir satisfying $\mathrm{ad}(L_{-1})(\mathrm{Vir}^-) \subseteq \mathrm{Vir}^-$. Therefore, $\mathrm{gr}(U(\mathrm{Vir}^-)) \cong \mathbb{C}[L_{-1}, L_{-2}, \ldots]$ has a derivation given by $\partial(L_{-n}) = (n-1)L_{-n-1}$ for $n \in \mathbb{Z}_+$.

Let Γ be an abelian group, and let $\mathfrak{g} = \bigoplus_{\alpha \in \Gamma} \mathfrak{g}^{\alpha}$ be a Γ -graded Lie algebra. A \mathfrak{g} -module $M = \bigoplus_{\alpha \in \Gamma} M^{\alpha}$ is Γ -graded if

$$\mathfrak{g}^{\alpha}M^{\beta} \subseteq M^{\alpha+\beta} \quad \text{for } \alpha, \beta \in \Gamma.$$

An \mathfrak{h} -module M is \mathfrak{h} -diagonalizable if

$$M = \bigoplus_{\lambda \in \mathfrak{h}^*} M_\lambda,$$

where

$$M_{\lambda} = \{ u \in M \mid \text{for } h \in \mathfrak{h}, \, hu = \lambda(h)u \}.$$

An \mathfrak{h} -diagonalizable module M is called \mathfrak{h} -semisimple if

$$\dim(M_{\lambda}) < \infty$$
 for $\lambda \in \mathfrak{h}^*$.

A $(\mathfrak{g}, \mathfrak{h})$ -module is an \mathfrak{h} -diagonalizable \mathfrak{g} -module. From now on, we assume that π_Q is injective. The character of a semisimple $(\mathfrak{g}, \mathfrak{h})$ -module M is given by

$$\operatorname{ch}_M(q) = \sum_{\lambda \in \mathfrak{h}^*} \dim(M_\lambda) q^\lambda$$

The category $\mathcal{C}_{(\mathfrak{g},\mathfrak{h})}$ is the category whose objects are \mathfrak{h}^* -graded $(\mathfrak{g},\mathfrak{h})$ -modules and for $M, N \in \mathcal{C}_{(\mathfrak{g},\mathfrak{h})}$ $\mathcal{C}_{(\mathfrak{g},\mathfrak{h})}, \operatorname{Hom}_{\mathcal{C}_{(\mathfrak{g},\mathfrak{h})}}(M,N) = \operatorname{Hom}_{\mathfrak{g}}(M,N)$ (Lie algebra module homomorphisms).

Remark 3.2. If $f \in \operatorname{Hom}_{\mathfrak{g}}(M, N)$, then f respects the \mathfrak{h}^* -grading automatically because for $u \in M_{\lambda}$ and $h \in \mathfrak{h}$, $hf(u) = f(hu) = f(\lambda(h)u) = \lambda(h)f(u)$.

We can introduce a partial order on Q as follows

 $\lambda_1 \leq \lambda_2 \iff$ there exists $\gamma \in Q^+$ such that $\lambda_2 - \lambda_1 = \pi_Q(\gamma)$.

Since π_Q is assumed to be injective, we can abuse notation and write $\lambda_2 - \lambda_1 = \gamma \in Q$. We now define the full subcategory $\mathcal{C}_{(\mathfrak{g},\mathfrak{h})}^{\lambda \geq}$ by considering the objects $M \in \mathcal{C}_{(\mathfrak{g},\mathfrak{h})}$ such that $M_{\mu} = 0$ if $\lambda \geq \mu$.

Example 3.3. When considering the \mathbb{Z} -graded Virasoro Lie algebra (Vir, \mathfrak{h}), where $\mathfrak{h} = \mathbb{C}L_0 \oplus \mathbb{C}C$, we identify \mathfrak{h}^* with \mathbb{C}^2 as $\lambda = (c, h)$ if $\lambda(C) = c$ and $\lambda(L_0) = h$. We are mainly interested in the highest weight modules M(c, h) and L(c, h) of the Virasoro Lie algebra. In this case, $(c_2, n_2) \geq$ (c_1, n_1) if and only if $c_1 = c_2$ and $n_1 - n_2 \in \mathbb{N}$. If this happens, $(c_2, n_2) - (c_1, n_1) = n_1 - n_2$.

4. The refined character

Let $(\mathfrak{g}, \mathfrak{h})$ be a \mathbb{Z} -graded Lie algebra, let $\lambda \in \mathfrak{h}^*$, and let $M \in \mathcal{C}_{(\mathfrak{g},\mathfrak{h})}^{\lambda \geq}$. The *PBW filtration of* M is given by

$$M^p = \{au \mid q \in \mathbb{N}, \mu \le \lambda, a \in U(\mathfrak{g}^-)^q, u \in M_\mu, q + \lambda - \mu \le p\} \text{ for } p \in \mathbb{Z}.$$

This filtration clearly satisfies axioms (i)–(iv) of §2 with $U(\mathfrak{g}^-)$ in place of A, and gr(M) becomes a gr $(U(\mathfrak{g}^{-}))$ -module.

If $f: M \to N$ is a homomorphism, where $M, N \in \mathcal{C}_{(\mathfrak{a}, \mathfrak{b})}^{\lambda \geq}$, then

$$\begin{split} &\operatorname{gr}(f):\operatorname{gr}(M)\to\operatorname{gr}(N),\\ &\operatorname{gr}(f)(\gamma^p_M(u))=\gamma^p_N(f(u))\quad\text{for }p\in\mathbb{N}\text{ and }u\in M^p \end{split}$$

defines a homomorphism of $\operatorname{gr}(U(\mathfrak{g}^{-}))$ -modules. Therefore, we obtain a functor

$$\operatorname{gr}: \mathcal{C}_{(\mathfrak{g},\mathfrak{h})}^{\lambda \geq} \to \operatorname{gr}(U(\mathfrak{g}^{-}))$$
-Mod.

Remark 4.1. Let $M, N \in \mathcal{C}_{(\mathfrak{g},\mathfrak{h})}^{\lambda \geq}$, and let $f: M \to N$ be a homomorphism. By Remark 3.2, f respects the \mathfrak{h}^* -grading, and this implies $f(M^p) \subseteq N^p$.

Example 4.2 (gr(M(c,h))). We pick a highest weight (c,h). By [Dix96, §2], we have a natural isomorphism

$$\operatorname{gr}(M(c,h)) \xrightarrow{\sim} \mathbb{C}[L_{-1}, L_{-2}, \dots] \cong \operatorname{gr}(U(\operatorname{Vir}^{-})),$$

 $\gamma^{\operatorname{len}(\lambda)}(L_{\lambda}|c,h\rangle) \mapsto p_{\lambda} \quad \text{for } \lambda \text{ a composition.}$

In this article, we are interested in the *refined character of* $M \in \mathcal{C}_{(\mathfrak{g},\mathfrak{h})}^{\lambda \geq}$, where M is semisimple, defined to be

$$\operatorname{ch}_M(t,q) = \sum_{p \in \mathbb{N}} \sum_{\lambda \in \mathfrak{h}^*} \dim(\gamma^p(M^p \cap M_\lambda)) t^p q^\lambda,$$

By axiom (iii) of A-modules given in §2, we have

$$\operatorname{ch}_M(1,q) = \operatorname{ch}_M(q).$$

Example 4.3 $(ch_{M(c,h)}(t,q))$. We consider the \mathbb{Z} -graded Virasoro Lie algebra $(Vir, \mathbb{C}L_0 \oplus \mathbb{C}C)$. For the Verma representation M(c,h), we have:

$$\operatorname{ch}_{M(c,h)}(t,q) = \frac{q^h}{\prod_{k \in \mathbb{Z}_+} (1 - tq^k)},$$
$$\operatorname{ch}_{M(c,h)}(q) = \frac{q^h}{\prod_{k \in \mathbb{Z}_+} (1 - q^k)} = \frac{q^h}{(q)_{\infty}}$$

Example 4.4 $(ch_{L(1/2,0)}(t,q))$. In [AVEH22], the refined character of the Virasoro minimal model Vir_{3,4} = L(1/2,0), also known as the *Ising model*, is computed explicitly. Interestingly, this character is connected to Nahm sums for the matrix $\begin{pmatrix} 8 & 3 \\ 3 & 2 \end{pmatrix}$ (see [Nah07]) and can be explicitly expressed as

$$\operatorname{ch}_{L(1/2,0)}(t,q) = \sum_{k_1,k_2 \in \mathbb{N}} t^{2k_1+k_2} \frac{q^{4k_1^2+3k_1k_2+k_2^2}}{(q)_{k_1}(q)_{k_2}} (1-q^{k_1}+q^{k_1+k_2}),$$

where $(q)_k = \prod_{j=1}^k (1-q^j) \in \mathbb{C}[q]$ denotes the *q*-Pochhammer symbol.

Remark 4.5. Let V be an N-graded conformal vertex algebra, and let M be an $h + \mathbb{N}$ -graded V-module. It is worth noting that the PBW filtration $(M^p)_{p \in \mathbb{Z}}$ and the standard filtration $(G^p M)_{p \in \mathbb{Z}}$, as introduced in [Sal24], are different. Therefore, the associated graded objects $\operatorname{gr}(M)$ and $\operatorname{gr}^G(M)$ and the refined characters $\operatorname{ch}_M(t,q)$ and $\operatorname{ch}_{\operatorname{gr}^G(M)}(t,q)$ are both different. For example, for a highest weight (c,h),

$$\operatorname{ch}_{\operatorname{gr}^{G}(M(c,h))}(t,q) = \frac{q^{h}}{(1-tq)\prod_{k\geq 2}(1-t^{2}q^{k})}$$

We pick a highest weight (c, h). We have a natural epimorphism

$$\pi_{c,h} : M(c,h) \twoheadrightarrow L(c,h),$$

$$\pi_{c,h}(u) = u + J(c,h),$$

and it satisfies $\ker(\pi_{c,h}) = J(c,h)$. Applying the functor gr, we obtain an epimorphism of modules over $\operatorname{gr}(\operatorname{Vir}^{-})$

$$\operatorname{gr}(\pi_{c,h}) : \operatorname{gr}(M(c,h)) \twoheadrightarrow \operatorname{gr}(L(c,h))$$

and this produces a natural isomorphism of modules over $gr(Vir^{-})$

$$\operatorname{gr}(M(c,h))/K(c,h) \xrightarrow{\sim} \operatorname{gr}(L(c,h)),$$

where

$$K(c,h) = \ker(\operatorname{gr}(\pi_{c,h})).$$

Explicitly, we have

$$K(c,h) = \sum_{p \in \mathbb{N}} \gamma^p (J(c,h) \cap M(c,h)^p).$$
(4.1)

From these observations, we obtain the following four isomorphisms which are going to be used frequently implicitly.

Proposition 4.6. We pick a highest weight (c, h). We have four (conformal) weight-preserving vector space isomorphisms:

$$\begin{split} M(c,h) &\xrightarrow{\sim} \operatorname{gr}(M(c,h)), \\ L_{\lambda}|c,h\rangle &\mapsto \gamma^{\operatorname{len}(\lambda)}(L_{\lambda}|c,h\rangle), \\ \operatorname{gr}(M(c,h)) &\xrightarrow{\sim} \mathbb{C}[L_{-1},L_{-2},\dots], \\ \gamma^{\operatorname{len}(\lambda)}(L_{\lambda}|c,h\rangle) &\mapsto p_{\lambda}, \\ L(c,h) &\xrightarrow{\sim} \operatorname{gr}(L(c,h)), \\ L_{\lambda}(|c,h\rangle + J(c,h)) &\mapsto \gamma^{\operatorname{len}(\lambda)}(L_{\lambda}(|c,h\rangle + J(c,h))), \\ \operatorname{gr}(L(c,h)) &\xrightarrow{\sim} \operatorname{gr}(M(c,h))/K(c,h), \\ \gamma^{\operatorname{len}(\lambda)}(L_{\lambda}(|c,h\rangle + J(c,h))) &\mapsto \gamma^{\operatorname{len}(\lambda)}(L_{\lambda}|c,h\rangle) + K(c,h), \end{split}$$

where λ is a partition.

In this article, we will deal with the polynomial algebra $\mathbb{C}[L_{-1}, L_{-2}, \ldots]$, its ideals and quotients. We will always use the degree reverse lexicographic order with $L_{-1} > L_{-2} > \ldots$. Actually, we can only use Gröbner basis theory with finitely generated polynomial algebras like $\mathbb{C}[L_{-1}, L_{-2}, \ldots, L_{-N}]$ for some $N \in \mathbb{N}$. But taking $N \to \infty$, we can obtain our desired results, see [Sal24] for details regarding this.

5. Fields over vector spaces

The vector space of formal distributions in $n \in \mathbb{N}$ variables, denoted by $\mathbb{C}[[x_1^{\pm 1}, \ldots, x_n^{\pm 1}]]$, is the set of functions $f : \mathbb{Z}^n \to \mathbb{C}$, written as $f(x_1, \ldots, x_n) = \sum_{m_1, \ldots, m_n \in \mathbb{Z}} f_{m_1, \ldots, m_n} x_1^{m_1} \ldots x_n^{m_n}$, with the natural operations of addition and multiplication by a scalar. The field of formal Laurent series, denoted by $\mathbb{C}((x))$, is the subspace of elements $f(x) \in \mathbb{C}[[x^{\pm 1}]]$ such that there is $N \in \mathbb{Z}$ with $f_n = 0$ for $n \leq N$. We also have $\mathbb{C}((x)) = \operatorname{Frac}(\mathbb{C}[[x]])$, so $\mathbb{C}((x))$ is actually a field. If V is a vector space, we similarly define $V[[x_1^{\pm 1}, \ldots, x_n^{\pm 1}]]$ and V((x)), but in this case, V((x)) is only a vector space.

Let V be a vector space. The Fourier expansion of a formal distribution $a(z) \in V[[z^{\pm 1}]]$, written as $a(z) = \sum_{n \in \mathbb{Z}} a_n z^n$, is conventionally written in the theory of vertex algebras as

$$a(z) = \sum_{n \in \mathbb{Z}} a_{(n)} z^{-n-1},$$

where

$$a_{(n)} = a_{-n-1}$$

Let V be a vector space, and let $a(z) \in \text{End}(V)[[z^{\pm 1}]]$ be a formal distribution. We set:

$$a(z)_{+} = \sum_{n \le -1} a_{(n)} z^{-n-1}$$
$$a(z)_{-} = \sum_{n \ge 0} a_{(n)} z^{-n-1}.$$

A formal distribution a(z) is a *field* if

$$a(z)b = \sum_{n \in \mathbb{Z}} a_{(n)}bz^{-n-1} \in V((z)) \quad \text{for } b \in V.$$

The vector space of fields over V is denoted by $\mathcal{F}(V)$. We note that

$$\mathcal{F}(V) = \operatorname{Hom}(V, V((z))).$$

Therefore, we can define a field a(z) by defining $a(z)b \in V((z))$ for $b \in V$.

Proposition 5.1 ([Noz08, Proposition 3.3.2]). Let $a(z), b(z) \in \mathcal{F}(V)$ be two fields. Then : $a(z)b(z) :\in \operatorname{End}(V)[[z^{\pm 1}]]$ is again a field, where :a(z)b(z) : is defined by

$$: a(z)b(z) : c = a(z)_{+}b(z)c + b(z)a(z)_{-}c \text{ for } c \in V.$$

We thus defined the notion of normal ordered product between fields $a(z), b(z) \in \mathcal{F}(V)$, denoted by : a(z)b(z) :. In general, the operation of normal ordered product is neither commutative nor associative. We follow the convention that the normal ordered product is read from right to left, so that, by definition,

$$a(z)b(z)c(z) :=: a(z)(:b(z)c(z):):$$

Lemma 5.2 ([Noz08, Proposition 3.3.3]). Let $a(z), b(z) \in \mathcal{F}(V)$ be two fields. Their normal ordered product is written explicitly as

$$: a(z)b(z) := \sum_{j \in \mathbb{Z}} : a(z)b(z) :_{(j)} z^{-j-1}$$

with

$$: a(z)b(z):_{(j)} c = \sum_{n \le -1} a_{(n)}b_{(j-n-1)}c + \sum_{n \ge 0} b_{(j-n-1)}a_{(n)}c \quad for \ c \in V.$$

Lemma 5.3. Let V be a vector space. We consider s fields $a^1(z), \ldots, a^s(z) \in \mathcal{F}(V)$, with $s \ge 2$, and let $b \in V$. For $l \in \mathbb{Z}$,

$$: a^{1}(z)a^{2}(z)\dots a^{s}(z):_{(l)} b = \sum_{n_{1},\dots,n_{s-1}\in\mathbb{N}}\sum_{k=0}^{s-1} R^{l,k}_{n_{1},\dots,n_{s-1}}(a^{1}(z),\dots,a^{s}(z))b,$$

where $R_{n_1,\dots,n_{s-1}}^{l,k}(a^1(z),\dots,a^s(z))$ is the sum of $\binom{s-1}{k}$ terms given by

$$R_{n_1,\dots,n_{s-1}}^{l,k}(a^1(z),\dots,a^s(z)) = \sum_{\substack{1 \le i_1 < \dots < i_k \le s-1 \\ 1 \le j_1 < \dots < j_{s-1-k} \le s-1 \\ \{i_1,\dots,i_k\} \cup \{j_1,\dots,j_{s-1-k}\} = \{1,\dots,s-1\}} a_{(-n_{j_1}-1)}^{j_1} \dots a_{(-n_{j_{s-1-k}}-1)}^{j_{s-1-k}} a_{(l-k-\sum_{r=1}^k n_{i_r} + \sum_{r=1}^{s-1-k} n_{j_r})} a_{(n_{i_k})}^{i_k} \dots a_{(n_{i_1})}^{i_1}.$$

Proof. This follows from Lemma 5.2 and induction on s.

6. Modules over the simple Virasoro vertex algebras

Let $c \in \mathbb{C}$. We make the subalgebra $\operatorname{Vir}^{\geq} \oplus \mathbb{C}L_{-1}$ of Vir act on \mathbb{C} as follows:

 $L_n 1 = 0$ for $n \ge -1$ and C1 = c.

The induced Vir-module

$$\operatorname{Vir}^{c} = \operatorname{Ind}_{\operatorname{Vir}^{\geq} \oplus \mathbb{C}L_{-1}}^{\operatorname{Vir}} = U(\operatorname{Vir}) \otimes_{U(\operatorname{Vir}^{\geq} \oplus \mathbb{C}L_{-1})} \mathbb{C}$$

carries the structure of a vertex algebra and is called the *universal Virasoro vertex algebra of* central charge c, see [FBZ01, §2] or [Kac17, §2] for details.

The vertex algebra Vir^c has a unique maximal proper ideal. We denote by Vir_c the unique simple quotient, and we call it the *simple Virasoro vertex algebra of central charge c*. Let $p, q \geq 2$ be relatively prime integers, and we set

$$c_{p,q} = 1 - \frac{6(p-q)^2}{pq}$$

Let V be a Vir-module. A vector u in V is called *singular* if it is nonzero and

$$L_n u = 0 \quad \text{for } n \in \mathbb{Z}_+.$$

Theorem 6.1 ([GK07]). *The following are equivalent:*

(i) Vir^{c} is not simple, i.e., $\operatorname{Vir}^{c} \neq \operatorname{Vir}_{c}$;

(ii) c is of the form $c_{p,q}$ for some $p,q \ge 2$ relatively prime integers.

Moreover, the maximal proper ideal of $\operatorname{Vir}^{c_{p,q}}$ is generated by a singular vector of conformal weight (p-1)(q-1), denoted by $a_{p,q}$. In the expression

$$a_{p,q} = \sum_{\substack{\lambda \vdash (p-1)(q-1)\\\lambda_{\text{len}(\lambda)} \ge 2}} c_{\lambda} L_{\lambda} |0\rangle,$$

where $c_{\lambda} \in \mathbb{Q}$, the coefficient of $L_{-2}^{(p-1)(q-1)/2}$ is nonzero.

Proposition 6.2. Let V be a vertex algebra, and let $Y^M : V \to \mathcal{F}(M)$ be a V-module. For $s, n_1, \ldots, n_s \in \mathbb{N}$ and $a^1, \ldots, a^s \in V$,

$$Y^{M}(a_{(-n_{1}-1)}^{1}\dots a_{(-n_{s}-1)}^{s}|0\rangle, z) = \frac{:\partial_{z}^{n_{1}}Y^{M}(a^{1}, z)\dots\partial_{z}^{n_{s}}Y^{M}(a^{s}, z):}{n_{1}!\dots n_{s}!}$$

Proof. This is a consequence of the *n*-product identity for modules over vertex algebras ([LL04, (5.2.16)]).

We are interested in the irreducible modules over the simple Virasoro vertex algebras $\operatorname{Vir}_{p,q} = \operatorname{Vir}_{c_{p,q}} = L(c_{p,q}, 0)$, where $p, q \geq 2$ are relatively prime integers. In this context, the following constants play a crucial role. For integers m, n such that 0 < m < p and 0 < n < q, we set

$$h_{m,n} = \frac{(np - mq)^2 - (p - q)^2}{4pq}$$

Theorem 6.3 ([Wan93]). We set $c = c_{p,q}$ for some $p,q \ge 2$ relatively prime integers. Then the irreducible modules over Vir_c are $L(c, h_{m,n})$ for integers m, n such that 0 < m < p and 0 < n < q. Let $Y_{\operatorname{Vir}^c}^{L(c,h_{m,n})}$: $\operatorname{Vir}^c \to \mathcal{F}(L(c, h_{m,n}))$ be the state-field correspondence of $L(c, h_{m,n})$ as a module over Vir^c , and let $Y_{\operatorname{Vir}^c}^{L(c,h_{m,n})}$: $\operatorname{Vir}_c \to \mathcal{F}(L(c, h_{m,n}))$ be the state-field correspondence of $L(c, h_{m,n})$ as a module over Vir_c . Then $Y_{\operatorname{Vir}^c}^{L(c,h_{m,n})}$ factors through $Y_{\operatorname{Vir}_c}^{L(c,h_{m,n})}$, i.e., the following diagram commutes

$$\begin{array}{c} \operatorname{Vir}^{c} & \longrightarrow & \operatorname{Vir}_{c} \\ & & & \downarrow_{Y_{\operatorname{Vir}^{c}}^{L(c,h_{m,n})}} \\ & & & \downarrow_{Y_{\operatorname{Vir}^{c}}^{L(c,h_{m,n})}} \\ & & \mathcal{F}(L(c,h_{m,n})) \end{array}$$

where the horizontal arrow is the quotient map.

We are also interested in the refined character of these irreducible modules. The ordinary character is fortunately already known.

Theorem 6.4 ([FF84]). Let $p, q \ge 2$ be relatively prime integers, and let m, n be integers such that 0 < m < p and 0 < n < q. Then

$$\operatorname{ch}_{L(c_{p,q},h_{m,n})}(q) = \frac{1}{(q)_{\infty}} \sum_{k \in \mathbb{Z}} q^{\frac{(2kpq+mq-np)^2 - (p-q)^2}{4pq}} - q^{\frac{(2kpq+mq+np)^2 - (p-q)^2}{4pq}}$$
$$= \frac{q^{h_{m,n}}}{(q)_{\infty}} \sum_{k \in \mathbb{Z}} q^{k^2 pq + k(mq-np)} - q^{k^2 pq + k(mq+np) + mn}.$$

In this article, we consider:

$$p = 2,$$

$$q = 2s + 1 \quad \text{for } s \in \mathbb{Z}_+.$$

By Theorem 6.3, $\operatorname{Vir}_{2,2s+1}$ has s irreducible modules $L(c_{2,2s+1}, h_{1,1}), \ldots, L(c_{2,2s+1}, h_{1,s})$, known collectively as the boundary minimal models.

7. The LI FILTRATION

Let V be a vertex algebra, and let $(a^i)_{i \in I}$ be a family of strong generators of V. For $p \in \mathbb{Z}$, we set

$$F_p V = \operatorname{span}\{a_{(-n_1-1)}^{i_1} \dots a_{(-n_s-1)}^{i_s} | 0 \rangle | s, n_1, \dots, n_s \in \mathbb{N}, i_1, \dots, i_s \in I, n_1 + \dots + n_s \ge p\}.$$

Proposition 7.1 ([Li05]). The filtration $(F_pV)_{p\in\mathbb{Z}}$ satisfies:

(i)
$$F_pV = V$$
 for $p \leq 0$;
(ii) $|0\rangle \in F_0V \supseteq F_1V \supseteq \ldots$;
(iii) $T(F_pV) \subseteq F_{p+1}V$ for $p \in \mathbb{Z}$;
(iv) $a_{(n)}F_qV \subseteq F_{p+q-n-1}V$ for $p, q \in \mathbb{Z}$, $a \in F_pV$ and $n \in \mathbb{Z}$;
(v) $a_{(n)}F_qV \subseteq F_{p+q-n}V$ for $p, q \in \mathbb{Z}$, $a \in F_pV$ and $n \in \mathbb{N}$.

Let

$$\operatorname{gr}_F(V) = \bigoplus_{p \in \mathbb{N}} F_p V / F_{p+1} V$$

be the associated graded vector space. By [Li05], the vector space $\operatorname{gr}_F(V)$ is a vertex Poisson algebra with operations given as follows. For $p, q \in \mathbb{N}$, $a \in F_pV$ and $b \in F_qV$, we set:

$$\sigma_p(a)\sigma_q(b) = \sigma_{p+q}(a_{(-1)}b),$$

$$T(\sigma_p(a)) = \sigma_{p+1}(Ta),$$

$$Y_-(\sigma_p(a), z)\sigma_q(b) = \sum_{n \in \mathbb{N}} \sigma_{p+q-n}(a_{(n)}b)z^{-n-1},$$

where $\sigma_p: F_pV \to \operatorname{gr}_F(V)$ is the *principal symbol map*, which is the composition of the natural maps $F_pV \twoheadrightarrow F_pV/F_{p+1}V$ and $F_pV/F_{p+1}V \hookrightarrow \operatorname{gr}_F(V)$. The unit is $\sigma^0(|0\rangle)$. The filtration $(F_pV)_{p\in\mathbb{Z}}$ is called the *Li filtration of* V.

Lemma 7.2 ([Li05, Lemma 2.9]). Let V be a vertex algebra. Then

$$F_p V = \operatorname{span}\{a_{(-i-1)}b \mid a \in V, i \in \mathbb{Z}_+, b \in F_{p-i}V\} \quad for \ p \in \mathbb{Z}_+.$$

By Lemma 7.2, the Li filtration depends only on V and not on the choice of the strong generators. If V is graded with Hamiltonian H, then $H(F_pV) \subseteq F_pV$ for $p \in \mathbb{Z}$. Therefore, we can define an operator $H \in \text{End}(\text{gr}_F(V))$ as $H(\sigma_p(a)) = \sigma_p(Ha)$ for $p \in \mathbb{N}$ and $a \in F_pV$. For $p \in \mathbb{Z}$ and $\Delta \in \mathbb{C}$, we define $F_pV_{\Delta} = F_pV \cap V_{\Delta}$. Since $H(F_pV) \subseteq F_pV$ for $p \in \mathbb{Z}$, we have

$$F_p V = \bigoplus_{\Delta \in \mathbb{C}} F_p V_\Delta \quad \text{for } p \in \mathbb{Z}.$$

For $\Delta \in \mathbb{C}$, we define $\operatorname{gr}_F(V)_{\Delta} = \bigoplus_{p \in \mathbb{N}} \sigma_p(F_p V_{\Delta})$. Then $Ha = \Delta a$ for $a \in \operatorname{gr}_F(V)_{\Delta}$. The family of subspaces $(\operatorname{gr}_F(V)_{\Delta})_{\Delta \in \mathbb{C}}$ satisfies $\operatorname{gr}_F(V) = \bigoplus_{\Delta \in \mathbb{C}} \operatorname{gr}_F(V)_{\Delta}$. Therefore, the operator $H \in \operatorname{End}(\operatorname{gr}_F(V))$ is diagonalizable with $\operatorname{gr}_F(V)_{\Delta} = \operatorname{ker}(H - \Delta \operatorname{Id}_{\operatorname{gr}_F(V)})$. In fact, it is possible to prove that this diagonalizable operator H is actually a Hamiltonian of $\operatorname{gr}_F(V)$.

We have the natural vector space isomorphisms

$$\sigma_p(F_pV_\Delta) \cong F_pV_\Delta/F_{p+1}V_\Delta \quad \text{for } p \in \mathbb{Z} \text{ and } \Delta \in \mathbb{C}$$

and the refined grading

$$\operatorname{gr}_{F}(V) = \bigoplus_{\substack{p \in \mathbb{N} \\ \Delta \in \mathbb{C}}} \sigma_{p}(F_{p}V_{\Delta}).$$
(7.1)

By (7.1), when dim $(V_{\Delta}) < \infty$ for $\Delta \in \mathbb{C}$, it is natural to define the refined character of V with respect to the Li filtration as

$$\operatorname{ch}_{\operatorname{gr}_F(V)}(t,q) = \sum_{\substack{p \in \mathbb{N} \\ \Delta \in \mathbb{C}}} \dim(\sigma_p(F_p V_\Delta)) t^p q^{\Delta}.$$

If $f: V_1 \to V_2$ is a homomorphism of vertex algebras, then

$$\operatorname{gr}_F(f) : \operatorname{gr}_F(V_1) \to \operatorname{gr}_F(V_2), \operatorname{gr}_F(f)(\sigma_p^{V_1}(a)) = \sigma_p^{V_2}(f(a)) \quad \text{for } p \in \mathbb{N} \text{ and } a \in F_p V_1$$

defines a homomorphism of vertex Poisson algebras. If V_1 and V_2 are graded, then we require that f respects the gradings of V_1 and V_2 , and this implies that $\operatorname{gr}_F(f)$ also respects the gradings of $\operatorname{gr}_F(V_1)$ and $\operatorname{gr}_F(V_2)$. Therefore, we obtain a functor

 $\operatorname{gr}_F : \{(\operatorname{graded}) \text{ vertex algebras}\} \to \{(\operatorname{graded}) \text{ vertex Poisson algebras}\}.$

Example 7.3 (gr_F(Vir^c)). We pick $c \in \mathbb{C}$. We have an isomorphism

$$\operatorname{gr}_{F}(\operatorname{Vir}^{c}) \xrightarrow{\sim} \mathbb{C}[L_{-2}, L_{-3}, \dots],$$

 $\sigma_{\Delta(\lambda)-\operatorname{clen}(\lambda)}(L_{\lambda}|0\rangle) \mapsto p_{\lambda} \quad \text{for } \lambda \text{ a composition without ones.}$

The derivation $T \in \text{Der}(\mathbb{C}[L_{-2}, L_{-3}, ...])$ is given by $T(L_{-n}) = (n-1)L_{-n-1}$ for $n \ge 2$, which is extended to a derivation. The Poisson structure of $\text{gr}^G(\text{Vir}^c)$ is trivial (i.e., the map Y_- is zero). The refined character is given explicitly by

$$ch_{{
m gr}_F({
m Vir}^c)}(t,q) = rac{1}{\prod_{k\geq 2}(1-t^{k-2}q^k)}.$$

Let V be a vertex algebra, let $(a^i)_{i \in I}$ be a family of strong generators of V, and let M be a V-module. For $p \in \mathbb{Z}$, we set

$$F_p M = \operatorname{span}\{a_{(-n_1-1)}^{i_1 M} \dots a_{(-n_s-1)}^{i_s M} u \mid s, n_1, \dots, n_s \in \mathbb{N}, i_1, \dots, i_s \in I, u \in M, n_1 + \dots + n_s \ge p\}.$$

Proposition 7.4 ([Li05]). The filtration $(F_pM)_{p\in\mathbb{Z}}$ satisfies:

- (i) $M = F_p M$ for $p \le 0$; (ii) $F_0 M \supseteq F_1 M \supseteq \dots$;
- (iii) $a_{(n)}F_qM \subseteq F_{p+q-n-1}M$ for $p,q \in \mathbb{Z}$, $a \in F_pV$ and $n \in \mathbb{Z}$;
- (iv) $a_{(n)}F_qM \subseteq F_{p+q-n}M$ for $p, q \in \mathbb{Z}$, $a \in F_pV$ and $n \in \mathbb{N}$.

Let

$$\operatorname{gr}_F(M) = \bigoplus_{p \in \mathbb{N}} F_p M / F_{p+1} M$$

be the associated graded vector space. By [Li05], the vector space $\operatorname{gr}_F(M)$ is a module over $\operatorname{gr}_F(V)$ with operations given as follows. For $p, q \in \mathbb{N}$, $a \in F_pV$ and $u \in F_qM$, we set:

$$\sigma_p(a)\sigma_q^M(u) = \sigma_{p+q}^M(a_{(-1)}^M u),$$

$$Y_{-}^M(\sigma_p(a), z)\sigma_q^M(u) = \sum_{n \in \mathbb{N}} \sigma_{p+q-n}^M(a_{(n)}^M u) z^{-n-1},$$

where $\sigma_p^M : F_p M \to \operatorname{gr}_F(M)$ is the principal symbol map. The filtration $(F_p M)_{p \in \mathbb{Z}}$ is called the Li filtration of M.

Lemma 7.5 ([Li05, Lemma 2.9]). Let V be a vertex algebra, and let M be a V-module. Then $F_pM = \operatorname{span}\{a^M_{(-i-1)}u \mid a \in V, i \in \mathbb{Z}_+, u \in F_{p-i}M\} \quad for \ p \in \mathbb{Z}_+.$

By Lemma 7.5, the Li filtration depends only on M and not on the choice of the strong generators of V. If V is graded with Hamiltonian H, and M is graded with Hamiltonian H^M , then $H^M(F_pM) \subseteq F_pM$ for $p \in \mathbb{Z}$. Therefore, we can define an operator $H^M \in \text{End}(\text{gr}_F(M))$ as $H^M(\sigma_p^M(u)) = \sigma_p^M(H^M u)$ for $p \in \mathbb{N}$ and $u \in F_pM$. For $p \in \mathbb{Z}$ and $\Delta \in \mathbb{C}$, we define $F_pM_\Delta = F_pM \cap M_\Delta$. Since $H^M(F_pM) \subseteq F_pM$ for $p \in \mathbb{Z}$, we have

$$F_p M = \bigoplus_{\Delta \in \mathbb{C}} F_p M_\Delta \quad \text{for } p \in \mathbb{Z}.$$

For $\Delta \in \mathbb{C}$, we define $\operatorname{gr}_F(M)_{\Delta} = \bigoplus_{p \in \mathbb{N}} \sigma_p^M(F_p M_{\Delta})$. Then $H^M u = \Delta u$ for $u \in \operatorname{gr}_F(M)_{\Delta}$. The family of subspaces $(\operatorname{gr}_F(M)_{\Delta})_{\Delta \in \mathbb{C}}$ satisfies $\operatorname{gr}_F(M) = \bigoplus_{\Delta \in \mathbb{C}} \operatorname{gr}_F(M)_{\Delta}$. Therefore, the operator $H^M \in \operatorname{End}(\operatorname{gr}_F(M))$ is diagonalizable with $\operatorname{gr}_F(M)_{\Delta} = \ker(H^M - \Delta \operatorname{Id}_{\operatorname{gr}_F(M)})$. In fact, it is possible to prove that this diagonalizable operator H^M is actually a Hamiltonian of $\operatorname{gr}_F(M)$.

We have the natural vector space isomorphisms

$$\sigma_p^M(F_pM_\Delta) \cong F_pM_\Delta/F_{p+1}M_\Delta \quad \text{for } p \in \mathbb{Z} \text{ and } \Delta \in \mathbb{C}$$

and the refined grading

$$\operatorname{gr}_{F}(M) = \bigoplus_{\substack{p \in \mathbb{N} \\ \Delta \in \mathbb{C}}} \sigma_{p}(F_{p}M_{\Delta}).$$
(7.2)

By (7.2), when $\dim(M)_{\Delta} < \infty$ for $\Delta \in \mathbb{C}$, it is natural to define the refined character of M with respect to the Li filtration as

$$\operatorname{ch}_{\operatorname{gr}_{F}(M)}(t,q) = \sum_{\substack{p \in \mathbb{N} \\ \Delta \in \mathbb{C}}} \dim(\sigma_{p}(F_{p}M_{\Delta}))t^{p}q^{\Delta}.$$

If $f: M_1 \to M_2$ is a homomorphism of V-modules, then

$$\operatorname{gr}_F(f) : \operatorname{gr}_F(M_1) \to \operatorname{gr}_F(M_2),$$

$$\operatorname{gr}_F(f)(\sigma_p^{M_1}(u)) = \sigma_p^{M_2}(f(u)) \quad \text{for } p \in \mathbb{N} \text{ and } u \in F_pM_1$$

defines a homomorphism of $\operatorname{gr}_F(V)$ -modules. If M_1 and M_2 are graded, then we require that f respects the gradings of M_1 and M_2 , and this implies that $\operatorname{gr}_F(f)$ also respects the gradings of $\operatorname{gr}_F(M_1)$ and $\operatorname{gr}_F(M_2)$. Therefore, we obtain a functor

$$\operatorname{gr}_F : \{(\operatorname{graded}) \ V \operatorname{-modules}\} \to \{(\operatorname{graded}) \ \operatorname{gr}_F(V) \operatorname{-modules}\}.$$

Example 7.6 $(\operatorname{gr}_F(M(c,h)))$. We pick a highest weight (c,h). From Example 7.3, $\operatorname{gr}_F(\operatorname{Vir}^c)$ is isomorphic to $\mathbb{C}[L_{-2}, L_{-3}, \ldots]$. We have an isomorphism

$$\operatorname{gr}_{F}(M(c,h)) \xrightarrow{\sim} \bigoplus_{k \in \mathbb{N}} \mathbb{C}[L_{-2}, L_{-3}, \dots] L_{-1}^{k},$$
$$\sigma_{\Delta(\lambda) - \operatorname{clen}(\lambda)}(L_{\lambda}|c,h\rangle) \mapsto u_{\lambda} \quad \text{for } \lambda \text{ a composition.}$$

The refined character is given explicitly by

$$\operatorname{ch}_{\operatorname{gr}_F(M(c,h))}(t,q) = \frac{q^n}{(1-q)\prod_{k\geq 2}(1-t^{k-2}q^k)}.$$

8. CLASSICALLY FREE VERTEX ALGEBRAS AND THEIR MODULES

Let V be a vertex algebra. By Lemma 7.2, we have

$$C_2V = \text{span}\{a_{(-2)}b \mid a, b \in V\} = F_1V.$$

We define the Zhu C_2 -algebra of V by

$$R_V = V/C_2 V = F_0 V/F_1 V \subseteq \operatorname{gr}_F(V).$$

In Zhu's original article [Zhu96] and other works, R_V is denoted by A(V).

The fact that $\operatorname{gr}_F(V)$ is a vertex Poisson algebra implies that R_V is a Poisson algebra with operations given as follows. For $a, b \in V$, we set:

$$\sigma_0(a)\sigma_0(b) = \sigma_0(a_{(-1)}b),$$

$$\{\sigma_0(a), \sigma_0(b)\} = \sigma_0(a_{(0)}b).$$

If V is graded, then, as we explained in §7, $\operatorname{gr}_F(V)$ is graded. Thus, R_V is also graded. We have constructed a functor

 $R: \{(\text{graded}) \text{ vertex algebras}\} \rightarrow \{(\text{graded}) \text{ Poisson algebras}\}.$

Oftentimes, some condition on R_V implies or is equivalent to some condition on V. The vertex algebra V is called C_2 -cofinite if R_V is finite dimensional.

Example 8.1 (R_{Vir^c} and R_{Vir_c}). We pick $c \in \mathbb{C}$. We consider $\mathbb{C}[L_{-2}]$ as the polynomial algebra in one variable L_{-2} , and we equip it with the trivial Poisson bracket. By Example 7.3, we have the following isomorphism of Poisson algebras

$$R_{\operatorname{Vir}^{c}} \xrightarrow{\sim} \mathbb{C}[L_{-2}],$$

$$\sigma_{0}(L_{-2} |0\rangle) \mapsto L_{-2}.$$

We now move onto Vir_c. If c is not of the form $c_{p,q}$ for some $p, q \ge 2$ relatively prime integers, then $\operatorname{Vir}_c = \operatorname{Vir}^c$ by Theorem 6.1, and we have already solved the problem. Therefore, we assume c is of this form.

We have a natural quotient map

$$\pi_c : \operatorname{Vir}^c \twoheadrightarrow \operatorname{Vir}_c,$$

$$\pi_c(a) = a + U(\operatorname{Vir})\{a_{p,q}\}$$

Applying the functor R, we obtain an epimorphism

$$R_{\pi_c}: R_{\operatorname{Vir}^c} \twoheadrightarrow R_{\operatorname{Vir}_c}.$$

From the equation $\ker(R_{\pi_c}) = \sigma_0(U(\operatorname{Vir})\{a_{p,q}\})$ and Theorem 6.1, we obtain

$$\ker(R_{\pi_c}) = (\sigma_0(L_{-2}^{(p-1)(q-1)/2} |0\rangle)).$$

In summary,

$$R_{\operatorname{Vir}_{p,q}} \cong \mathbb{C}[L_{-2}]/(L_{-2}^{(p-1)(q-1)/2})$$

Thus, Vir^c is never C_2 -cofinite, while Vir_c is C_2 -cofinite only when c is of the form $c_{p,q}$ for some $p, q \ge 2$ relatively prime integers.

Lemma 8.2 ([Li05, Corollary 4.3]). Let V be a vertex algebra. As a differential algebra, $gr_F(V)$ is generated by R_V , i.e.,

$$\operatorname{gr}_F(V) = (R_V)_T.$$

Let V be a (graded) vertex algebra. We have a natural (graded) algebra inclusion inc : $R_V \hookrightarrow \operatorname{gr}_F(V)$. By the universal property of inc : $R_V \hookrightarrow JR_V$ (see Appendix B), there is a (graded) differential algebra homomorphism $\phi_V : JR_V \to \operatorname{gr}_F(V)$ such that the following diagram commutes



Because R_V is a (graded) Poisson algebra, we can equip JR_V with the level 0 vertex Poisson algebra structure, as explained in Appendix C. From now on, JR_V will be considered as a (graded) vertex Poisson algebra.

Lemma 8.3. Let V be a (graded) vertex algebra. The (graded) differential algebra homomorphism $\phi_V : JR_V \to \operatorname{gr}_F(V)$ defined above is surjective and is actually a (graded) vertex Poisson algebra homomorphism.

Proof. The homomorphism ϕ_V is surjective by Lemma 8.2. The fact that ϕ_V is a (graded) vertex Poisson algebra homomorphism is explained in [Ara12, Proposition 2.5.1].

When ϕ_V is an isomorphism, we say V is classically free.

Example 8.4 (JR_{Vir^c} and JR_{Vir_c}). Let $c \in \mathbb{C}$. Then Vir^c is classically free because

$$\operatorname{gr}_F(\operatorname{Vir}^c) \cong \mathbb{C}[L_{-2}, L_{-3}, \dots]$$

by Example 7.3 and

$$JR_{\operatorname{Vir}^c} \cong J(C[L_{-2}]) = \mathbb{C}[L_{-2}, L_{-3}, \dots]$$

by Example 8.1 and Appendix B.

We now move onto Vir_c. If c is not of the form $c_{p,q}$ for some $p, q \ge 2$ relatively prime integers, then $\operatorname{Vir}_c = \operatorname{Vir}^c$ by Theorem 6.1, and we have already solved the problem. Therefore, we assume c is of this form. Then

$$JR_{\operatorname{Vir}_{p,q}} \cong J(\mathbb{C}[L_{-2}]/(L_{-2}^{(p-1)(q-1)/2})) = \mathbb{C}[L_{-2}, L_{-3}, \dots]/(L_{-2}^{(p-1)(q-1)/2})_{\delta}$$

by Example 8.1 and Appendix B.

Example 8.5. If $q > p \ge 2$ are relatively prime integers, then $\operatorname{Vir}_{p,q}$ is classically free if and only if p = 2 by [VEH21].

Example 8.6. It was proven in [AVEH22] that the Ising model $Vir_{3,4}$ is not classically free. In fact, by [AVEH22, Theorem 2],

$$\ker(\phi_{\mathrm{Vir}_{3,4}}) = (b)_{\partial_2}$$

where

$$b = L_{-4}L_{-3}L_{-2} + \frac{1}{6}L_{-5}L_{-2}^2$$

and $(b)_{\partial}$ is the differential ideal generated by b, cf. Appendix B and Example 7.3 where ∂ is denoted by T.

Let V be a vertex algebra, and let M be a module over V. By Lemma 7.5, we have

$$C_2M = \operatorname{span}\{a_{(-2)}^M \mid a \in V, u \in M\} = F_1M.$$

We define the Zhu C_2 -module of M by

$$R_M = M/C_2M = F_0M/F_1M \subseteq \operatorname{gr}_F(M).$$

The fact that $\operatorname{gr}_F(M)$ is a module over $\operatorname{gr}_F(V)$ implies that R_M is a module over R_V with operations given as follows. For $a \in V$ and $u \in M$, we set:

$$\sigma_0(a)\sigma_0^M(u) = \sigma_0^M(a_{(-1)}^M u)$$

$$\{\sigma_0(a), \sigma_0^M(u)\} = \sigma_0^M(a_{(0)}^M u).$$

If V is graded and M is a graded V-module, then, as we explained in §7, $\operatorname{gr}_F(M)$ is a graded $\operatorname{gr}_F(V)$ -module. Thus, R_M is also a graded R_V -module.

We have constructed a functor

$$R: \{(\text{graded}) \ V\text{-modules}\} \rightarrow \{(\text{graded}) \ R_V\text{-modules}\}$$

Example 8.7 $(R_{M(c,h)} \text{ and } R_{L(c,h)})$. We pick a highest weight (c, h). As in Example 8.1, we consider $\mathbb{C}[L_{-2}]$ as the polynomial algebra in one variable L_{-2} , and we equip it with the trivial Poisson bracket. We consider $\bigoplus_{k \in \mathbb{N}} \mathbb{C}[L_{-2}]L_{-1}^k \cong \mathbb{C}[L_{-2}, L_{-1}]$ as a module over the Poisson algebra $\mathbb{C}[L_{-2}]$ with Poisson bracket given by $\{L_{-2}, L_{-1}^k\} = L_{-1}^{k+1}$ for $k \in \mathbb{N}$. By Example 7.6 and Example 8.1, we have the following isomorphism of modules over Poisson algebras

$$R_{M(c,h)} \xrightarrow{\sim} \bigoplus_{k \in \mathbb{N}} \mathbb{C}[L_{-2}]L_{-1}^{k}$$
$$\sigma_{0}(L_{-2}|c,h\rangle) \mapsto L_{-2},$$
$$\sigma_{0}(L_{-1}|c,h\rangle) \mapsto L_{-1}.$$

As in Example 8.1, we obtain

$$R_{L(c,h)} \cong \frac{\bigoplus_{k \in \mathbb{N}} \mathbb{C}[L_{-2}]L_{-1}^k}{\sigma_0(J(c,h))}.$$

Theorem 8.8 ([Li05, Lemma 4.2]). Let V be a vertex algebra, and let M be a V-module. As a $\operatorname{gr}_F(V)$ -module without the Poisson structure, $\operatorname{gr}_F(M)$ is generated by R_M , i.e.,

$$\operatorname{gr}_F(M) = \operatorname{gr}_F(V)R_M.$$

Let V be a (graded) vertex algebra, and let M be a (graded) V-module. We have a natural (graded) inclusion inc : $R_M \hookrightarrow \operatorname{gr}_F(M)$ of (graded) R_V -modules. As we explained in Appendix C, we can consider $JR_V \otimes_{R_V} R_M$ as a (graded) JR_V -module. Because we have a (graded) epimorphism of vertex Poisson algebras $\phi_V : JR_V \twoheadrightarrow \operatorname{gr}_F(V)$, we can consider the (graded) $\operatorname{gr}_F(V)$ -module $\operatorname{gr}_F(M)$ as a (graded) JR_V -module. By the universal property of inc : $R_M \hookrightarrow JR_V \otimes_{R_V} R_M$ (see Appendix C), there is a (graded) JR_V -module homomorphism $\phi_M : JR_V \otimes_{R_V} R_M \to \operatorname{gr}_F(M)$ such that the following diagram commutes

$$\begin{array}{ccc} R_M & \xleftarrow{\operatorname{inc}} & JR_V \otimes_{R_V} R_M \\ & & & & \downarrow \phi_M \\ & & & & \operatorname{gr}_F(M) \end{array}$$

Lemma 8.9. The (graded) JR_V -module homomorphism $\phi_M : JR_V \otimes_{R_V} R_M \to \operatorname{gr}_F(M)$ defined above is surjective.

Proof. The assertion follows from Lemma 8.3 and Theorem 8.8.

When ϕ_M is an isomorphism, we say M is *classically free*.

Example 8.10 $(JR_{\text{Vir}^c} \otimes_{R_{\text{Vir}^c}} R_{M(c,h)})$. We pick a highest weight (c, h). The Verma module M(c, h) is always classically free as a Vir^c-module because by Example 7.6, Example 8.1 and Example 8.7,

$$JR_{\operatorname{Vir}^{c}} \otimes_{R_{\operatorname{Vir}^{c}}} R_{M(c,h)} \cong \mathbb{C}[L_{-2}, L_{-3}, \dots] \otimes_{\mathbb{C}[L_{-2}]} \bigoplus_{k \in \mathbb{N}} \mathbb{C}[L_{-2}]L_{-1}^{k}$$
$$\cong \bigoplus_{k \in \mathbb{N}} \mathbb{C}[L_{-2}, L_{-3}, \dots]L_{-1}^{k}$$
$$\cong \operatorname{gr}_{F}(M(c, h)).$$

We wish to determine when the irreducible modules $L(c_{p,q}, h_{m,n})$ over the simple Virasoro vertex algebras $\operatorname{Vir}_{p,q}$ are classically free, as mentioned in Theorem 0.3 and Theorem 6.3.

9. Proofs of the main theorems

In this section, we fix $s \in \mathbb{Z}_+$ and $i \in \{1, \ldots, s\}$.

A partition $\lambda = [\lambda_1, \ldots, \lambda_m]$ contains a partition $\eta = [\eta_1, \ldots, \eta_n]$, written as $\eta \subseteq \lambda$, if $m \ge n$ and there is $i \in \mathbb{Z}_+$ such that $1 \le i \le m - n + 1$ and $[\lambda_i, \lambda_{i+1}, \ldots, \lambda_{i+n-1}] = \eta$.

We define

$$p^{s,i}(t,q) = \sum_{\lambda \in P^{s,i}} t^{\operatorname{len}(\lambda)} q^{\Delta(\lambda)} \in \mathbb{C}[[t,q]],$$

where $P^{s,i}$ is the set of partitions that do not contain any partition in $R^{s,i}$ as defined in §0, i.e.,

$$P^{s,i} = \{ \lambda \vdash \mid \text{for } \eta \in R^{s,i}, \lambda \not\supseteq \eta \}$$

Lemma 9.1. We have

$$p^{s,i}(t,q) = \sum_{k=(k_1,\dots,k_{s-1})\in\mathbb{N}^{s-1}} t^{kB_{s-1}^{(s)}} \frac{q^{\frac{1}{2}kG^{(s)}k^T + kB_{s-i}^{(s)}}}{(q)_{k_1}\dots(q)_{k_{s-1}}}.$$

Proof. Following the notation of [And98, §7], we write:

$$p^{s,i}(t,q) = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} b_{s,i}(m,n) t^m q^n,$$
$$J_{s,i}(0,t,q) = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} c_{s,i}(m,n) t^m q^n.$$

By [And98, §7.3], we have $b_{s,i}(m,n) = c_{s,i}(m,n)$ for $m, n \in \mathbb{N}$. This means $p^{s,i}(t,q) = J_{s,i}(0,t,q)$. By the proof of [And98, Theorem 7.8], we have

$$J_{s,i}(0,t,q) = \sum_{k=(k_1,\dots,k_{s-1})\in\mathbb{N}^{s-1}} \frac{t^{N_1+\dots+N_{s-1}}q^{N_1^2+\dots+N_{s-1}^2+N_i+\dots+N_{s-1}}}{(q)_{k_1}\dots(q)_{k_{s-1}}},$$

where $N_j = k_j + \cdots + k_{s-1}$ for $j = 1, \ldots, s$. Since:

$$N_1^2 + \dots + N_{s-1}^2 = \frac{1}{2}kG^{(s)}k^T,$$

 $N_j + \dots + N_{s-1} = kB_{s-j}^{(s)}$ for $j = 1, \dots, s_s$

we have

$$\sum_{k=(k_1,\dots,k_{s-1})\in\mathbb{N}^{s-1}}\frac{t^{N_1+\dots+N_{s-1}}q^{N_1^2+\dots+N_{s-1}^2+N_i+\dots+N_{s-1}}}{(q)_{k_1}\dots(q)_{k_{s-1}}} = \sum_{k=(k_1,\dots,k_{s-1})\in\mathbb{N}^{s-1}}t^{kB_{s-1}^{(s)}}\frac{q^{\frac{1}{2}kG^{(s)}k^T+kB_{s-i}^{(s)}}}{(q)_{k_1}\dots(q)_{k_{s-1}}}.$$

Lemma 9.2. We have

$$\operatorname{ch}_{L(c_{2,2s+1},h_{1,i})}(q) = q^{h_{1,i}} p^{s,i}(1,q) = q^{h_{1,i}} \left(\prod_{\substack{n \neq 0, \pm i \mod 2s+1}}^{\infty} (1-q^n)^{-1}\right).$$

Proof. By the proof of Lemma 9.1, [And98, Lemma 7.3] and [And98, Corollary 2.9], we have

$$\begin{aligned} q^{h_{1,i}} p^{s,i}(1,q) &= q^{h_{1,i}} J_{s,i}(0,1,q) \\ &= q^{h_{1,i}} \left(\prod_{\substack{n \neq 0, \pm i \text{ mod } 2s+1}}^{\infty} (1-q^n)^{-1} \right) \\ &= \frac{q^{h_{1,i}}}{(q)_{\infty}} \sum_{k \in \mathbb{N}} (-1)^k q^{(2s+1)k(k+1)/2-ki} (1-q^{(2k+1)i}) \\ &= \frac{q^{h_{1,i}}}{(q)_{\infty}} \sum_{k \in \mathbb{Z}} (-1)^k q^{(2s+1)k(k+1)/2-ki}. \end{aligned}$$

The disjoint union $\mathbb{Z} = \{2k \mid k \in \mathbb{Z}\} \cup \{-2k - 1 \mid k \in \mathbb{Z}\}$ and Theorem 6.4 imply that

$$\begin{aligned} \frac{q^{h_{1,i}}}{(q)_{\infty}} \sum_{k \in \mathbb{Z}} (-1)^k q^{(2s+1)k(k+1)/2-ki} &= \frac{q^{h_{1,i}}}{(q)_{\infty}} \left(\sum_{k \in \mathbb{Z}} (-1)^{2k} q^{(2s+1)2k(2k+1)/2-2ki} \right. \\ &\quad + \sum_{k \in \mathbb{Z}} (-1)^{-2k-1} q^{(2s+1)(-2k-1)(-2k-1+1)/2-(-2k-1)i} \right) \\ &\quad = \frac{q^{h_{1,i}}}{(q)_{\infty}} \left(\sum_{k \in \mathbb{Z}} q^{2k^2(2s+1)+k(2s+1-2i)} - \sum_{k \in \mathbb{Z}} q^{2k^2(2s+1)+k(2s+2i+1)+i} \right) \\ &\quad = \mathrm{ch}_{L(c_{2,2s+1},h_{1,i})}(q). \end{aligned}$$

We order the PBW basis of $U(\text{Vir}^-) = \text{span}\{L_{\lambda} \mid \lambda \vdash\}$ by degree reverse lexicographic order with $L_{-1} > L_{-2} > \ldots$. Formally, for any partitions λ and η , we define

 $L_{\lambda} \leq L_{\eta}$ if and only if $p_{\lambda} \leq p_{\eta}$.

For $x \in U(\text{Vir}^-)$ with $x \neq 0$, we may write

$$x = c_1 L_{\lambda^1} + c_2 L_{\lambda^2} + \dots + c_r L_{\lambda^r},$$

where for $1 \leq i \leq r, 0 \neq c_i \in \mathbb{C}$ and $L_{\lambda^1} > L_{\lambda^2} > \cdots > L_{\lambda^r}$. We define the *leading power of* x as $lp(x) = L_{\lambda^1}$. We set lp(0) = 0. Next, for a highest weight (c, h), we extend the definition of lp from $U(\text{Vir}^-)$ to M(c, h) by considering the isomorphism of vector spaces $U(\text{Vir}^-) \xrightarrow{\sim} M(c, h), L_{\lambda} \mapsto L_{\lambda} | c, h \rangle$, where λ is a partition.

Remark 9.3. The definition of the order in the PBW basis of $U(\text{Vir}^-)$ was made so that for a highest weight (c, h), a partition λ and $u \in M(c, h)$, if $\ln(u) = L_{\lambda}|c, h\rangle$, then $\ln(\gamma^{\ln(\lambda)}(u)) = p_{\lambda}$.

Lemma 9.4. For $\lambda \in \mathbb{R}^{s,i}$, there exists $u \in J(c_{2,2s+1}, h_{1,i})$ homogeneous such that $lp(u) = L_{\lambda}|c_{2,2s+1}, h_{1,i}\rangle$.

Proof. By Theorem 6.1, the maximal proper ideal of $\operatorname{Vir}^{c_{2,2s+1}}$ is generated by a singular vector of conformal weight 2s, denoted by $a_{2,2s+1}$, which has the form

$$a_{2,2s+1} = L_{-2}^{s} |0\rangle + \sum_{\substack{\mu \vdash 2s \\ \mu_{\ln(\mu)} \ge 2 \\ \ln(\mu) < s}} c_{\mu} L_{\mu} |0\rangle,$$
(9.1)

where $c_{\mu} \in \mathbb{Q}$. We have two cases:

(i) $\lambda = [a^d, (a+1)^{s-d}]$ for some $a \in \mathbb{Z}_+$ and $d \in \{1, \ldots, s\}$. We set

$$u = (a_{2,2s+1})_{(s-sa+d-1)} | c_{2,2s+1}, h_{1,i} \rangle$$

and claim

$$lp(u) = L_{\lambda} | c_{2,2s+1}, h_{1,i} \rangle.$$

To prove this, we first prove that

$$(L_{-2}^{s}|0\rangle)_{(s-sa+d-1)}|c_{2,2s+1},h_{1,i}\rangle = {\binom{s}{d}}L_{\lambda}|c_{2,2s+1},h_{1,i}\rangle + \sum_{\substack{\mu\vdash\Delta(\lambda)\\ \ln(\mu)\leq \ln(\lambda)\\\mu\neq\lambda}}c_{\mu}L_{\mu}|c_{2,2s+1},h_{1,i}\rangle, \quad (9.2)$$

where $c_{\mu} \in \mathbb{Q}$. Applying Proposition 6.2 and Lemma 5.3 with:

$$a^{1}(z) = \dots = a^{s}(z) = L(z) = \sum_{n \in \mathbb{Z}} L_{(n)} z^{-n-1} = \sum_{n \in \mathbb{Z}} L_{n} z^{-n-2},$$

$$b = |c_{2,2s+1}, h_{1,i}\rangle,$$

$$l = s - sa + d - 1,$$

we obtain

$$(L_{-2}^{s}|0\rangle)_{(l)}|c_{2,2s+1},h_{1,i}\rangle = \sum_{n_{1},\dots,n_{s-1}\in\mathbb{N}}\sum_{k=0}^{s-1}R_{n_{1},\dots,n_{s-1}}^{l,k}(L(z),\dots,L(z))|c_{2,2s+1},h_{1,i}\rangle,$$
(9.3)

where $R_{n_1,\dots,n_{s-1}}^{l,k}(L(z),\dots,L(z))$ is the sum of $\binom{s-1}{k}$ terms given by

$$R_{n_1,\dots,n_{s-1}}^{l,k}(L(z),\dots,L(z)) = \sum_{\substack{1 \le i_1 < \dots < i_k \le s-1 \\ 1 \le j_1 < \dots < j_{s-1-k} \le s-1 \\ \{i_1,\dots,i_k\} \cup \{j_1,\dots,j_{s-1-k}\} = \{1,\dots,s-1\}} L_{(-n_{j_1}-1)} \dots L_{(-n_{j_{s-1-k}}-1)} L_{(l-k-\sum_{r=1}^k n_{i_r}+\sum_{r=1}^{s-1-k} n_{j_r})} L_{(n_{i_k})} \dots L_{(n_{i_1})},$$

where

$$L_{(-n_{j_1}-1)} \dots L_{(-n_{j_{s-1-k}}-1)} L_{(l-k-\sum_{r=1}^k n_{i_r}+\sum_{r=1}^{s-1-k} n_{j_r})} L_{(n_{i_k})} \dots L_{(n_{i_1})} = L_{-n_{j_1}-2} \dots L_{-n_{j_{s-1-k}}-2} L_{s-sa+d-2-k-\sum_{r=1}^k n_{i_r}+\sum_{r=1}^{s-1-k} n_{j_r}} L_{n_{i_k}-1} \dots L_{n_{i_1}-1}.$$

We see that each term L_{μ} appearing in $R_{n_1,\dots,n_{s-1}}^{l,k}(L(z),\dots,L(z))$, where μ is a combination (not necessarily a partition), satisfies $\Delta(\mu) = sa + s - d = \Delta(\lambda)$ and $\operatorname{len}(\mu) \leq \operatorname{len}(\lambda)$.

We recall that $L_{\lambda} - L_{\lambda\sigma} \in U(\text{Vir}^{-})^{\text{len}(\lambda)-1}$ for a combination λ and a permutation $\sigma \in S_{\text{len}(\lambda)}$ (see [Dix96, §2] for details). Therefore, for a combination μ , we can expand $L_{\mu}|_{c_{2,2s+1}}, h_{1,i}\rangle$ in the following way

$$L_{\mu}|c_{2,2s+1},h_{1,i}\rangle = L_{\text{prt}(\mu)}|c_{2,2s+1},h_{1,i}\rangle + \sum_{\substack{\mu\vdash\Delta(\lambda)\\ \text{len}(\mu)<\text{len}(\lambda)}} c_{\mu}L_{\mu}|c_{2,2s+1},h_{1,i}\rangle,$$

where $c_{\mu} \in \mathbb{Q}$. We need to compute the coefficient of $L_{\lambda}|c_{2,2s+1}, h_{1,i}\rangle$ in u when expressed as a linear combination of the elements of the usual PBW basis. To do this, we need to see how much each sum $R_{n_1,\dots,n_{s-1}}^{l,k}(L(z),\dots,L(z))|c_{2,2s+1},h_{1,i}\rangle$ contributes to the coefficient of $L_{\lambda}|c_{2,2s+1},h_{1,i}\rangle$ in u. We have two subcases:

Subcase a = 1: In the sum (9.3), if k > d or k < d-1 or $n_i \neq 0$ for some $i = 1, \ldots, s-1$, then $R_{n_1,\ldots,n_{s-1}}^{l,k}(L(z),\ldots,L(z))|c_{2,2s+1},h_{1,i}\rangle$ contributes 0. As for the two remaining sums, the sum $R_{0,\ldots,0}^{l,d-1}(L(z),\ldots,L(z))|c_{2,2s+1},h_{1,i}\rangle$ contributes $\binom{s-1}{d-1}$ because we are picking d-1 elements from $\{1,\ldots,s-1\}$, and the sum $R_{0,\ldots,0}^{l,d}(L(z),\ldots,L(z))|c_{2,2s+1},h_{1,i}\rangle$ contributes $\binom{s-1}{d}$ because we are picking d elements from $\{1,\ldots,s-1\}$. Since $\binom{s-1}{d-1} + \binom{s-1}{d} = \binom{s}{d}$, we obtain (9.2).

Subcase a > 1: In the sum (9.3), if k > 0, then $R_{n_1,\dots,n_{s-1}}^{l,k}(L(z),\dots,L(z))|_{c_{2,2s+1}},h_{1,i}\rangle$ contributes 0. We have

$$R_{n_1,\dots,n_{s-1}}^{l,0}(L(z),\dots,L(z)) = L_{-n_1-2}\dots L_{-n_{s-1}-2}L_{s-sa+d-2+\sum_{r=1}^{s-1}n_r}$$

Therefore, the sum $R_{n_1,\ldots,n_{s-1}}^{l,0}(L(z),\ldots,L(z))|_{c_{2,2s+1}},h_{1,i}\rangle$ contributes $\binom{s-1}{d-1}$ when exactly d-1 of the n_i s are equal to a-2 and s-d of the n_i s are equal to a-1, contributes $\binom{s-1}{d}$ when exactly d of the n_i s are equal to a-2 and s-d-1 of the n_i s are equal to a-1, and contributes 0 otherwise. Since $\binom{s-1}{d-1} + \binom{s-1}{d} = \binom{s}{d}$, we obtain (9.2).

(ii) $\lambda = [1^i]$. This follows from [BSA88], where it is proved that one of the generating singular vectors in $J(c_{2,2s+1}, h_{1,i})$ contains the term $L^i_{-1}|c_{2,2s+1}, h_{1,i}\rangle$ in its expansion with respect to the usual PBW basis.

We note that if μ is a partition of $\Delta(\lambda)$ such that $\operatorname{len}(\mu) \leq \operatorname{len}(\lambda)$ and $\mu \neq \lambda$, then $L_{\lambda} > L_{\mu}$. Thus, (9.2) implies $\operatorname{lp}((L_{-2}^{s}|0\rangle)_{(s-sa+d-1)}|c_{2,2s+1},h_{1,i}\rangle) = L_{\lambda}|c_{2,2s+1},h_{1,i}\rangle$. From (9.1) and Lemma 5.3, we see that $\operatorname{lp}(u) = \operatorname{lp}((L_{-2}^{s}|0\rangle)_{(s-sa+d-1)}|c_{2,2s+1},h_{1,i}\rangle) = L_{\lambda}|c_{2,2s+1},h_{1,i}\rangle$. Finally, by Theorem 6.3, $u \in J(c_{2,2s+1},h_{1,i})$, and it is clear that u is homogeneous from its definition. \Box

Remark 9.5. For any partitions λ and η , if $\lambda \supseteq \eta$, then $p_{\eta} \mid p_{\lambda}$. The converse is not true. For example, $p_{[4,2]} \mid p_{[4,3,2]}$, but $[4,3,2] \not\supseteq [4,2]$. However, if $\eta = [\eta_1, \ldots, \eta_m]$ and $\eta_1 - \eta_m \le 1$, then $\lambda \supseteq \eta$ if and only if $p_{\eta} \mid p_{\lambda}$.

Lemma 9.6. There is an alternative description for $P^{s,i}$, namely

$$P^{s,i} = \{ \lambda \vdash | \text{ for } \eta \in R^{s,i}, p_\eta \nmid p_\lambda \}.$$

Proof. This is a consequence of Remark 9.5.

Sketch of proof of Theorem 0.2. Let G be a Gröbner basis of $K(c_{2,2s+1}, h_{1,i})$, and we set:

$$B = \{ p_{\lambda} \mid \text{for } a \in G, \, \operatorname{lp}(a) \nmid p_{\lambda} \},\$$
$$\overline{B} = \{ p_{\lambda} \mid \lambda \in P^{s,i} \}.$$

By Gröbner basis theory, (4.1), Remark 9.3, Lemma 9.4 and Lemma 9.6, we have $\overline{B} \subseteq B$. Actually, Proposition 4.6 and Lemma 9.2 imply $\overline{B} = B$. We have found a basis of $\operatorname{gr}(L(c_{2,2s+1}, h_{1,i}))$ and by Proposition 4.6, a basis of $L(c_{2,2s+1}, h_{1,i})$. For details, see [Sal24].

Proof of Theorem 0.1. This is a corollary of Theorem 0.2, see [Sal24] for details.

Proof of Theorem 0.3. By Example 8.7, we have

$$R_{L(c_{2,2s+1},h_{1,i})} \cong \frac{\bigoplus_{k \in \mathbb{N}} \mathbb{C}[L_{-2}]L_{-1}^k}{\sigma_0(J(c_{2,2s+1},h_{1,i}))} \cong \frac{\mathbb{C}[L_{-2},L_{-1}]}{\sigma_0(J(c_{2,2s+1},h_{1,i}))}$$

First, we prove that for a partition η with $\eta = \emptyset$ or $2 \ge \eta_1$,

$$p_{\eta} + \sigma_0(J(c_{2,2s+1}, h_{1,i})) \in \text{span}\{p_{\lambda} + \sigma_0(J(c_{2,2s+1}, h_{1,i})) \mid \lambda = \emptyset \text{ or } (\lambda \in P^{s,i} \text{ and } 2 \ge \lambda_1)\} = W.$$

We assume, for the sake of contradiction, that there is a power p_{η} with $\eta = \emptyset$ or $2 \ge \eta_1$ not satisfying $p_{\eta} + \sigma_0(J(c_{2,2s+1}, h_{1,i})) \in W$.

Let p_{η_0} be the smallest of such powers in the degree reverse lexicographic order with $L_{-1} > L_{-2} > \ldots$. We must have $\eta_0 \neq \emptyset$ and $\eta_0 \notin P^{s,i}$, otherwise $p_{\eta_0} + \sigma_0(J(c_{2,2s+1}, h_{1,i})) \in W$. By Lemma 9.6, there is $\mu \in \mathbb{R}^{s,i}$ such that $p_{\mu} \mid p_{\eta_0}$. From (4.1), Remark 9.3, Lemma 9.4 and the fact that lp is multiplicative, we obtain $u \in J(c_{2,2s+1}, h_{1,i})$ homogeneous such that $p(u) = L_{\eta_0} | c_{2,2s+1}, h_{1,i} \rangle$.

From the equation

$$u = L_{\eta_0} | c_{2,2s+1}, h_{1,i} \rangle + \sum_{\substack{\mu \vdash \Delta(\eta_0) \\ L_{\mu} < L_{\eta_0}}} c_{\mu} L_{\mu} | c_{2,2s+1}, h_{1,i} \rangle,$$

where $c_{\mu} \in \mathbb{Q}$, we obtain

$$p_{\eta_0} + \sigma_0(J(c_{2,2s+1}, h_{1,i})) = -\sum_{\substack{\mu \vdash \Delta(\eta_0) \\ p_\mu < p_{\eta_0} \\ 2 \ge \mu_1}} c_\mu(p_\mu + \sigma_0(J(c_{2,2s+1}, h_{1,i}))).$$

The last equation implies $p_{\eta_0} + \sigma_0(J(c_{2,2s+1}, h_{1,i})) \in W$ because $p_\mu + \sigma_0(J(c_{2,2s+1}, h_{1,i})) \in W$ if $p_\mu < p_{\eta_0}$ by the minimality of p_{η_0} , and this is a contradiction. We have proved that

$$\frac{\mathbb{C}[L_{-2}, L_{-1}]}{\sigma_0(J(c_{2,2s+1}, h_{1,i}))} = \operatorname{span}\{p_\lambda + \sigma_0(J(c_{2,2s+1}, h_{1,i})) \mid \lambda = \emptyset \text{ or } (\lambda \in P^{s,i} \text{ and } 2 \ge \lambda_1)\}.$$
(9.4)

By [VEH21] or [BMS13], we already know $\operatorname{Vir}_{2,2s+1}$ is classically free. This means that $\operatorname{gr}_F(\operatorname{Vir}_{2,2s+1}) \cong JR_{\operatorname{Vir}_{2,2s+1}} \cong J(\mathbb{C}[L_{-2}]/(L_{-2}^s)) = \mathbb{C}[L_{-2}, L_{-3}, \dots]/(L_{-2}^s)_{\partial}$ and

$$\mathbb{C}[L_{-2}, L_{-3}, \dots]/(L_{-2}^s)_{\partial} = \bigoplus_{\lambda \in P^{s,1}} \mathbb{C}(p_{\lambda} + (L_{-2}^s)_{\partial}).$$
(9.5)

We have

$$JR_{\text{Vir}_{2,2s+1}} \otimes_{R_{\text{Vir}_{2,2s+1}}} R_{L(c_{2,2s+1},h_{1,i})} \cong \frac{\mathbb{C}[L_{-2},L_{-3},\dots]}{(L_{-2}^{s})_{\partial}} \otimes_{\frac{\mathbb{C}[L_{-2}]}{(L_{-2}^{s})}} \frac{\mathbb{C}[L_{-2},L_{-1}]}{\sigma_{0}(J(c_{2,2s+1},h_{1,i}))}.$$
(9.6)

Given a partition $\lambda \in P^{s,i}$, we can write λ uniquely as $\lambda = [\lambda^1, \lambda^2]$, where $\lambda^1 \in P^{s,1}$ satisfies $\lambda^1 = \emptyset$ or $\lambda^1_{\text{len}(\lambda^1)} \ge 3$ and $\lambda^2 = \emptyset$ or $(\lambda^2 \in P^{s,i} \text{ and } 2 \ge \lambda_1^2)$.

Given $\lambda \in P^{s,i}$, we define

$$t_{\lambda} = (p_{\lambda^{1}} + (L_{-2}^{s})_{\partial}) \otimes (p_{\lambda^{2}} + \sigma_{0}(J(c_{2,2s+1}, h_{1,i}))) \in \frac{\mathbb{C}[L_{-2}, L_{-3}, \dots]}{(L_{-2}^{s})_{\partial}} \otimes_{\frac{\mathbb{C}[L_{-2}]}{(L_{-2}^{s})}} \frac{\mathbb{C}[L_{-2}, L_{-1}]}{\sigma_{0}(J(c_{2,2s+1}, h_{1,i}))}.$$

From (9.4), (9.5) and (9.6), we conclude that

$$JR_{\text{Vir}_{2,2s+1}} \otimes_{R_{\text{Vir}_{2,2s+1}}} R_{L(c_{2,2s+1},h_{1,i})} = \text{span}\{t_{\lambda} \mid \lambda \in P^{s,i}\}.$$
(9.7)

By Theorem 0.2 and Proposition 4.6 with gr_F instead of gr , the natural epimorphism $\phi_{L(c_{2,2s+1},h_{1,i})}$ maps the set $\{t_{\lambda} \mid \lambda \in P^{s,i}\}$ onto a basis of $\operatorname{gr}_F(L(c_{2,2s+1},h_{1,i}))$. Thus, the set $\{t_{\lambda} \mid \lambda \in P^{s,i}\}$ is

linearly independent, and by (9.7), it is a basis of $JR_{\text{Vir}_{2,2s+1}} \otimes_{R_{\text{Vir}_{2,2s+1}}} R_{L(c_{2,2s+1},h_{1,i})}$. Therefore, $\phi_{L(c_{2,2s+1},h_{1,i})}$ is an isomorphism, and $L(c_{2,2s+1},h_{1,i})$ is classically free.

Let p, q > 2 be relatively prime integers, and let m, n be integers such that 0 < m < p and 0 < n < q. We have two natural inclusions inc : $JR_{\operatorname{Vir}_{p,q}} \hookrightarrow JR_{\operatorname{Vir}_{p,q}} \otimes_{R_{\operatorname{Vir}_{p,q}}} R_{L(c_{p,q},h_{m,n})}$, $\operatorname{inc}(a) = a \otimes 1$ and $\operatorname{inc} : \operatorname{gr}_F(\operatorname{Vir}_{p,q}) \hookrightarrow \operatorname{gr}_F(L(c_{p,q},h_{m,n}))$, $\operatorname{inc}(a) = a1$. We assume, for the sake of contradiction, that $\phi_{L(c_{p,q},h_{m,n})}$ is an isomorphism. This would imply that $\phi_{\operatorname{Vir}_{p,q}}$ is also an isomorphism, a contradiction to Example 8.5. Finally, $\phi_{L(c_{p,q},h_{m,n})}$ is not an isomorphism, and $L(c_{p,q},h_{m,n})$ is not classically free as a module over $\operatorname{Vir}_{p,q}$.

As in the proof of Lemma 9.4, let $w \in J(c_{2,2s+1}, h_{1,i})$ be the singular vector such that $lp(w) = L_{-1}^i | c_{2,2s+1}, h_{1,i} \rangle$, and we set:

$$u_k = \sigma_0((a_{2,2s+1})_{(k-1)} | c_{2,2s+1}, h_{1,i} \rangle) \quad \text{for } k = 1, \dots, i-1,$$

$$u_i = \sigma_0(w).$$

Then $u_1, \ldots, u_i \in \sigma_0(J(c_{2,2s+1}, h_{1,i}))$. Moreover:

$$L_{-2}^{s} = \sigma_{0}(a_{2,2s+1})\sigma_{0}(|c_{2,2s+1}, h_{1,i}\rangle) = \sigma_{0}((a_{2,2s+1})_{(-1)}|c_{2,2s+1}, h_{1,i}\rangle) \in \sigma_{0}(J(c_{2,2s+1}, h_{1,i})) = l_{-2}^{s-k}L_{-1}^{k} \quad \text{for } k = 1, \dots, i-1,$$
$$lp(u_{i}) = L_{-1}^{i}.$$

Corollary 9.7. With the notation above, we have:

$$\sigma_0(J(c_{2,2s+1}, h_{1,i})) = (L^s_{-2}, u_1, \dots, u_i)_{\text{psn}},$$
(9.8)

$$R_{L(c_{2,2s+1},h_{1,i})} \cong \frac{\bigoplus_{k \in \mathbb{N}} \mathbb{C}[L_{-2}]L_{-1}^k}{(L_{-2}^s, u_1, \dots, u_i)_{\text{psn}}},$$
(9.9)

$$\operatorname{gr}_{F}(L(c_{2,2s+1}, h_{1,i})) \cong \frac{\mathbb{C}[L_{-2}, L_{-3}, \dots]}{(L_{-2}^{s})_{\partial}} \otimes_{\frac{\mathbb{C}[L_{-2}]}{(L_{-2}^{s})}} \frac{\bigoplus_{k \in \mathbb{N}} \mathbb{C}[L_{-2}] L_{-1}^{k}}{(L_{-2}^{s}, u_{1}, \dots, u_{i})_{\operatorname{psn}}},$$
(9.10)

where psn denotes the Poisson submodule generated by the given subset.

Proof. We use the notation of the proof of Theorem 0.3. Since $\{t_{\lambda} \mid \lambda \in P^{s,i}\}$ is a basis of $JR_{\text{Vir}_{2,2s+1}} \otimes_{R_{\text{Vir}_{2,2s+1}}} R_{L(c_{2,2s+1},h_{1,i})}$, the set in (9.4) is a basis of $R_{L(c_{2,2s+1},h_{1,i})}$. We note that for a partition $\lambda \in R^{s,i}$ with $2 \geq \lambda_1$, there is $u \in (L^s_{-2}, u_1, \ldots, u_i)_{\text{psn}}$ such that $\ln(u) = u_{\lambda}$. Therefore, we can prove (9.8) similarly to how we proved (9.4). The isomorphisms (9.9) and (9.10) follow.

Remark 9.8. We have

$$u_1 = \sigma_0((a_{2,2s+1})_{(0)}|c_{2,2s+1}, h_{1,i}\rangle) = \{\sigma_0(a_{2,2s+1}), \sigma_0(|c_{2,2s+1}, h_{1,i}\rangle)\} = \{L_{-2}^s, 1\} = sL_{-2}^{s-1}L_{-1}.$$

However, I think there is no simple expression for u_2, \ldots, u_i and $R_{L(c_{2,2s+1},h_{1,i})}$.

APPENDIX A. GRADED POISSON ALGEBRAS AND THEIR MODULES

Let A be a commutative associative algebra. A Hamiltonian operator of A is a diagonalizable operator $H \in \text{End}(A)$ such that

$$H(ab) = H(a)b + aH(b)$$
 for $a, b \in A$.

Thus, a Hamiltonian of A is just a derivation of A. An algebra with a Hamiltonian is called *graded*.

Let A be a differential commutative associative algebra with derivation ∂ . A Hamiltonian operator of A is a Hamiltonian of A as a commutative associative algebra such that

$$[H,\partial] = \partial$$

It is possible to prove inductively that the last equation implies

$$H\partial^n = n\partial^n + \partial^n H \quad \text{for } n \in \mathbb{N}.$$
(A.1)

Let A be a Poisson algebra as defined in [Car03]. A Hamiltonian operator of A is a Hamiltonian H of A as a commutative associative algebra such that

$$H(\{a,b\}) = \{a, H(b)\} + \{H(a), b\} - \{a, b\} \text{ for } a, b \in A.$$

Let A be a graded commutative associative algebra with Hamiltonian H, and let M be an A-module. If A has a unit 1, we further assume that 1u = u for $u \in M$. A Hamiltonian operator of M is a diagonalizable operator $H^M \in \text{End}(M)$ such that

$$H^M(au) = H(a)u + aH^M(u)$$
 for $a \in A$ and $u \in M$.

Let A be a Poisson algebra. A module over A is an A-module M in the usual associative sense equipped with a bilinear map $\{\bullet, \bullet\} : A \times M \to M$, which makes M a Lie algebra module over A such that for $a, b \in A$ and $u \in M$:

- (i) (Left Leibniz rule) $\{a, bu\} = \{a, b\}u + b\{a, u\};$
- (ii) (Right Leibniz rule) $\{ab, u\} = a\{b, u\} + b\{a, u\}$.

Let A be a graded Poisson algebra with Hamiltonian H, and let M be an A-module. A Hamiltonian operator of M is a Hamiltonian H^M of M as a module over A as a commutative associative algebra such that

$$H^{M}(\{a,u\}) = \{a, H^{M}(u)\} + \{H(a), u\} - \{a, u\} \text{ for } a \in A \text{ and } u \in M.$$

Appendix B. Graded Jet Algebras and their modules

In this appendix, by an algebra we will mean a commutative associative algebra with unit. Let R be a finitely generated algebra. We now construct a differential algebra JR called the *jet algebra of* R and an algebra inclusion inc : $R \hookrightarrow JR$ universal with this property, i.e., for a differential algebra A and an algebra homomorphism $\phi : R \to A$, there is a unique differential algebra homomorphism $\overline{\phi} : JR \to A$ such that the following diagram commutes



Assuming $R = \mathbb{C}[x^1, \ldots, x^r]/(f_1, \ldots, f_s)$ for some polynomials $f_1, \ldots, f_s \in \mathbb{C}[x^1, \ldots, x^r]$, the construction is as follows. We introduce new variables $x_{(-i)}^j$ for $i = 1, 2, \ldots, j = 1, \ldots, r$ and a derivation ∂ of the polynomial algebra $\mathbb{C}[x_{(-i)}^j \mid i = 1, 2, \ldots, j = 1, \ldots, r]$ by setting

$$\partial x_{(-i)}^j = i x_{(-i-1)}^j$$
 for $i = 1, 2, \dots, j = 1, \dots, r$.

We set (identifying x^j with $x_{(-1)}^j$ when considering f_i in the following equation)

$$IR = \mathbb{C}[x_{(-i)}^{j} \mid i = 1, 2, \dots, j = 1, \dots, r] / (\partial^{j} f_{i} \mid i = 1, \dots, s, j = 0, 1, \dots)$$

= $\mathbb{C}[x_{(-i)}^{j} \mid i = 1, 2, \dots, j = 1, \dots, r] / (f_{1}, \dots, f_{s})_{\partial},$

where the subscript ∂ indicates the differential subalgebra generated by the given subset. By our definitions, ∂ factors through a derivation of JR, and we have an algebra inclusion inc : $R \hookrightarrow JR$, $\operatorname{inc}(x^j + (f_1, \ldots, f_s)) = x_{(-1)}^j + (f_1, \ldots, f_s)_\partial$ for $j = 1, \ldots, r$. The fact that inc : $R \hookrightarrow JR$ satisfies our desired universal property is explained in [Ara12, §2.3] and [EM09].

Remark B.1. We see that the classes of the original variables x^j generate JR as a differential algebra, i.e.,

$$JR = (x^1 + (f_1, \dots, f_s)_\partial, \dots, x^r + (f_1, \dots, f_s)_\partial)_\partial$$

If $f: R_1 \to R_2$ is a homomorphism of finitely generated algebras, then $Jf: JR_1 \to JR_2$ is defined by requiring that $Jf(\partial_1^n(\operatorname{inc}_1(x))) = \partial_2^n(\operatorname{inc}_2(f(x)))$ for $x \in R_1$ and $n \in \mathbb{N}$ (cf. Remark B.1). For a finitely generated algebra R and a differential algebra A, we have a natural isomorphism

 $\operatorname{Hom}_{\{\text{differential algebras}\}}(JR, A) \cong \operatorname{Hom}_{\{\text{algebras}\}}(R, A).$

Remark B.2. It is not true that we have defined a functor J which is left adjoint to the forgetful functor {differential algebras} \rightarrow {finitely generated algebras} because a differential algebra is generally not finitely generated.

But we can easily work in the general case as follows. Let R be an algebra (not necessarily finitely generated), and we consider the polynomial algebra in R variables $\mathbb{C}[(x^j)_{j\in R}]$. We have a natural epimorphism $\pi : \mathbb{C}[(x^j)_{j\in R}] \twoheadrightarrow R, \pi(x^j) = j$ for $j \in R$. We can repeat the construction of JR with $\mathbb{C}[(x^j)_{j\in R}]$ in place of $\mathbb{C}[x^1, \ldots, x^r]$ and ker (π) in place of (f_1, \ldots, f_s) . This way, we construct a functor

 $J: \{ algebras \} \to \{ differential \ algebras \},\$

which is left adjoint to the forgetful functor {differential algebras} \rightarrow {algebras}.

Given an algebra R, it is useful to consider the functor

 $JR \otimes_R \bullet : R\text{-Mod} \to JR\text{-Mod},$

where JR is merely considered an algebra. Again, the JR-module M together with the R-module inclusion inc : $M \hookrightarrow JR \otimes_R M$, $\operatorname{inc}(u) = 1 \otimes u$ satisfy a universal property similar to that of inc : $R \hookrightarrow JR$, and the functor $JR \otimes_R \bullet$ is left adjoint to the forgetful functor JR-Mod $\to R$ -Mod.

Let R be a graded algebra with Hamiltonian H. We can extend uniquely the Hamiltonian H to a Hamiltonian $H^{JR} \in \text{End}(JR)$ because of (A.1). Furthermore, if M is a graded R-module with Hamiltonian H^M , we can define a Hamiltonian of $JR \otimes_R M$ by setting

$$H^{JR\otimes_R M} = H^{JR} \otimes \mathrm{Id}_M + \mathrm{Id}_{JR} \otimes H^M.$$

APPENDIX C. GRADED VERTEX POISSON ALGEBRAS AND THEIR MODULES

Proposition C.1 ([Ara12, Proposition 2.3.1] and Appendix B). Let R be a (graded) Poisson algebra. Then there is a unique (graded) vertex Poisson algebra structure on JR such that

$$a_{(n)}b = \delta_{n,0}\{a, b\}$$
 for $a, b \in \mathbb{R}$ and $n \in \mathbb{N}$.

The (graded) vertex Poisson algebra structure on JR given in Proposition C.1 for a (graded) Poisson algebra R will be called the *level 0 vertex Poisson algebra structure of JR*.

If R is a (graded) Poisson algebra, and M is a (graded) R-module, then we can verify that $JR \otimes_R M$ is a (graded) JR-module by defining the Poisson structure as

$$a_{(n)}(b \otimes u) = (a_{(n)}b) \otimes u + \delta_{n,0}b \otimes \{a, u\}$$
 for $a \in R, b \in JR, u \in M$ and $n \in \mathbb{N}$

and $H^{JR\otimes_R M}$ as in Appendix B.

The (graded) *R*-module *M* together with the natural (graded) *R*-module inclusion inc : $M \hookrightarrow JR \otimes_R M$ satisfy the following universal property. Let *N* be a (graded) module over the (graded) vertex Poisson algebra JR, and let $f : M \to N$ be a (graded) homomorphism of modules over the (graded) Poisson algebra *R*. There is a unique (graded) homomorphism $\overline{f} : JR \otimes_R M \to N$ of (graded) modules over the (graded) vertex Poisson algebra *R* such that the following diagram commutes



References

- [And98] George E. Andrews. *The theory of partitions*. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 1998. Reprint of the 1976 original. (Cited on pages 2 and 17.)
- [Ara12] Tomoyuki Arakawa. A remark on the C_2 -cofiniteness condition on vertex algebras. Mathematische Zeitschrift, 270(1-2):559–575, February 2012. (Cited on pages 1, 14, 22, and 23.)
- [AVEH22] George E. Andrews, Jethro Van Ekeren, and Reimundo Heluani. The Singular Support of the Ising Model. International Mathematics Research Notices, page rnab328, May 2022. (Cited on pages 7 and 15.)

DIEGO SALAZAR

- [BMS13] Clemens Bruschek, Hussein Mourtada, and Jan Schepers. Arc spaces and the Rogers–Ramanujan identities. *The Ramanujan Journal*, 30(1):9–38, January 2013. (Cited on page 20.)
- [BSA88] Louis Benoit and Yvan Saint-Aubin. Degenerate conformal field theories and explicit expressions for some null vectors. *Physics Letters B*, 215(3):517–522, December 1988. (Cited on page 19.)
- [Car03] Paolo Caressa. Examples of Poisson modules, I. Rendiconti del Circolo Matematico di Palermo, 52(3):419–452, October 2003. (Cited on page 22.)
- [Dix96] Jacques Dixmier. *Enveloping algebras*, volume 11 of *Graduate studies in mathematics*. American Mathematical Society, Providence, R.I, 1996. (Cited on pages 4, 7, and 19.)
- [EM09] Lawrence Ein and Mircea Mustață. Jet schemes and singularities. In Algebraic geometry—Seattle 2005. Part 2, volume 80 of Proc. Sympos. Pure Math., pages 505–546. Amer. Math. Soc., Providence, RI, 2009. (Cited on page 22.)
- [FBZ01] Edward Frenkel and David Ben-Zvi. Vertex algebras and algebraic curves, volume 88 of Mathematical surveys and monographs. American Mathematical Society, Providence, R.I, 2001. (Cited on page 9.)
- [FF84] B. L. Feigin and D. B. Fuchs. Verma modules over the virasoro algebra. Functional Analysis and Its Applications, 17(3):241–242, 1984. (Cited on page 10.)
- [GK07] Maria Gorelik and Victor Kac. On simplicity of vacuum modules. Advances in Mathematics, 211(2):621–677, June 2007. (Cited on page 9.)
- [Gor61] Basil Gordon. A Combinatorial Generalization of the Rogers-Ramanujan Identities. American Journal of Mathematics, 83(2):393, April 1961. (Cited on page 1.)
- [IK11] Kenji Iohara and Yoshiyuki Koga. Representation theory of the Virasoro algebra. Springer Monographs in Mathematics. Springer-Verlag London, Ltd., London, 2011. (Cited on page 5.)
- [Kac17] Victor Kac. Introduction to Vertex Algebras, Poisson Vertex Algebras, and Integrable Hamiltonian PDE. In Filippo Callegaro, Giovanna Carnovale, Fabrizio Caselli, Corrado De Concini, and Alberto De Sole, editors, *Perspectives in Lie Theory*, volume 19, pages 3–72. Springer International Publishing, Cham, 2017. (Cited on page 9.)
- [KRR13] Victor G. Kac, A. K. Raina, and Natasha Rozhkovskaya. Bombay lectures on highest weight representations of infinite dimensional lie algebras, volume 29 of Advanced series in mathematical physics. World Scientific, Hackensack, New Jersey, Second edition, 2013. OCLC: ocn858312870. (Cited on page 3.)
- [Li05] Haisheng Li. Abelianizing Vertex Algebras. Communications in Mathematical Physics, 259(2):391–411, October 2005. (Cited on pages 1, 11, 12, 14, and 16.)
- [LL04] James Lepowsky and Haisheng Li. Introduction to Vertex Operator Algebras and Their Representations. Birkhäuser Boston, Boston, MA, 2004. (Cited on page 10.)
- [Nah07] Werner Nahm. Conformal Field Theory and Torsion Elements of the Bloch Group, pages 67–132. Springer Berlin Heidelberg, Berlin, Heidelberg, 2007. (Cited on page 7.)
- [Noz08] Christophe Nozaradan. Introduction to Vertex Algebras, November 2008. arXiv:0809.1380 [math-ph] version: 3. (Cited on page 9.)
- [Sal24] Diego Salazar. PBW bases of irreducible Ising modules. *Journal of Algebra*, 639:398–421, February 2024. (Cited on pages 7, 8, and 20.)
- [VEH21] Jethro Van Ekeren and Reimundo Heluani. Chiral Homology of Elliptic Curves and the Zhu Algebra. Communications in Mathematical Physics, 386(1):495–550, August 2021. (Cited on pages 1, 15, and 20.)
- [Wan93] Weiqiang Wang. Rationality of Virasoro vertex operator algebras. International Mathematics Research Notices, 1993(7):197–211, 1993. (Cited on page 10.)
- [Zhu96] Yongchang Zhu. Modular invariance of characters of vertex operator algebras. Journal of the American Mathematical Society, 9(1):237–302, 1996. (Cited on pages 1 and 13.)

INSTITUTO DE MATEMÁTICA PURA E APLICADA, RIO DE JANEIRO, RJ, BRAZIL Email address: diego.salazar@impa.br