### AUTOMORPHISM GROUPS OF CERTAIN ORBIFOLD VERTEX OPERATOR ALGEBRAS ARISING FROM COINVARIANT LATTICES ASSOCIATED WITH THE LEECH LATTICE

TAKARA KONDO

ABSTRACT. We determine the automorphism groups of the orbifold vertex operator algebras associated with the coinvariant lattices of isometries of the Leech lattice in the conjugacy classes 3C, 5C, 11A and 23A. These orbifold vertex operator algebras appear in a classification given by C.H. Lam and H. Shimakura.

### 1. INTRODUCTION

The orbifold of a vertex operator algebra (VOA) for an automorphism group is the fixed-points subVOA. We have a question: Are all automorphisms of its orbifold VOA obtained from the original VOA? Let us explain it precisely. For a VOA V and an automorphism group G of V,  $V^G$  is defined by the set of all fixed-points. Let V be simple and let G be a cyclic subgroup generated by a finite order automorphism of V. We have a group homomorphism from the normalizer  $N_{\text{Aut}(V)}(G)$  to  $\text{Aut}(V^G)$ . The main question is if the automorphism group  $Aut(V^G)$  is isomorphic to a quotient group of  $N_{\text{Aut}(V)}(G)$  or not. An automorphism of  $V^G$  is called *extra* if it cannot be obtained from the image of the normalizer  $N_{\text{Aut}(V)}(G)$ . It is important to consider when the orbifold VOA has extra automorphisms.

The lattice VOA  $V_L$  with respect to a positive-definite even lattice L is a significant example of VOA. Let g be a fixed-point free isometry of L. Then g can be lifted to an element  $\hat{g}$  of Aut  $(V_L)$ . When L has no root, the automorphism group Aut  $(V_L^{\hat{g}})$  is a finite group. We focus on such cases. In this paper, we consider the group structures of the automorphism groups of such orbifold VOAs.

The case that the order of g is 2, that is, g is the -1-isometry, has been studied in [Sh04];  $V_L^{\hat{g}}$  has extra automorphisms if and only if L can be constructed by Construction B from a doubly even binary code.

In [LS24], the case that g is a fixed-point free isometry of odd prime order p was treated and Lam and Shimakura classified all rootless even lattices such that  $V_L^{\hat{g}}$  has extra automorphisms. As to the above, Lam and Shimakura also classified rootless even lattices L such that the  $\tau$ -conjugate  $V_L(1) \circ \tau$  is of twisted type for some  $\tau \in \operatorname{Aut}(V_L^{\hat{g}})$ , where  $V_L(1)$  is the module defined in (3.2) (see Definition 2.4 for the definition of  $V_L(1)\circ\tau$ ). The following is the classification.

**Theorem 1.1.** ([LS24, Main Corollary 1]) Let L be a positive-definite rootless even lattice. Let g be a fixed-point free isometry of L of prime order p and let  $\hat{g}$  be a standard lift of g. Then the following are equivalent:

- There exists an automorphism  $\tau$  of  $V_L^{\hat{g}}$  such that  $V_L(1) \circ \tau$  is of twisted type;
- L is isometric to the coinvariant lattice  $\Lambda_{pX}$  of the Leech lattice  $\Lambda$  associated with the conjugacy class  $pX \in \{2A, -2A, 3B, 3C, 5B, 5C, 7B, 11A, 23A\}$ .

In [CLS18, La22+, Sh04], the automorphism groups associated with 2A, -2A, 3B, 5B and 7B were determined. So far the others have not been determined.

In this paper, we determine all of the remaining automorphism groups of the orbifold VOAs  $V_{\Lambda_{pX}}^{\hat{g}}$ , where  $pX \in \{3C, 5C, 11A, 23A\}$ . To determine the automorphism groups, we prove a generalization of [Sh04, Lemma 3.7](Theorem 5.1):

**Main Theorem 1.** Let L be a positive-definite rootless even lattice and let p be a prime number. Let g be a fixed-point free isometry of L. If  $(1 - g)L^* \subset L$ , then we have the following exact sequence

$$1 \longrightarrow A \longrightarrow \operatorname{Ker} \mu \stackrel{\tilde{\varphi}}{\longrightarrow} B \longrightarrow 1,$$

where  $A = \text{Hom}(L/(1-g)L^*, \mathbb{Z}_p)$ ,  $B = \{h \in C_{O(L)}(g) \mid h = 1 \text{ on } L^*/L\}/\langle g \rangle$  and the group homomorphism  $\mu$  is as in (5.1).

The following theorem describes the group structures of the remaining automorphism groups of the orbifold VOAs.

**Main Theorem 2.** Let  $(g, L) \in \{(3C, \Lambda_{3C}), (5C, \Lambda_{5C}), (11A, \Lambda_{11A}), (23A, \Lambda_{23A})\}$  and let  $\hat{g}$  be a lift of g as in Subsection 2.4. Then the automorphism groups of orbifold VOAs  $V_L^{\hat{g}}$  are following:

- Aut  $(V_{\Lambda_{3C}}^{\hat{g}}) \cong (3^4 \cdot (3^4 : 2)) \cdot (\Omega_7(3) \cdot 2));$
- Aut  $(V_{\Lambda_{5C}}^{g}) \cong (5^2 . (5^2 : 2)) . (2 \times \Omega_5(5));$
- Aut  $(V_{\Lambda_{11A}}^{\hat{g}}) \cong \Omega_4^-(11)$ . 2;
- Aut  $(V_{\Lambda_{23A}}^{\hat{g}}) \cong \Omega_3(23)$ . 2.

### 2. Preliminaries

2.1. Lattices. A *lattice* means a free abelian group of finite rank with a rational valued, positive-definite symmetric bilinear form  $(\cdot|\cdot)$ . Let L be a lattice with a positive-definite bilinear form  $(\cdot|\cdot)$ . The symbol O(L) denotes the isometry group of L. The *dual lattice* 

is defined by  $L^* = \{ \alpha \in \mathbb{Q} \otimes_{\mathbb{Z}} L \mid (\alpha | L) \subset \mathbb{Z} \}$ . A lattice is called *integral* if  $(\alpha | \beta) \in \mathbb{Z}$  for any  $\alpha, \beta \in L$ . A lattice is said to be *even* if  $(\alpha | \alpha) \in 2\mathbb{Z}$  for all  $\alpha \in L$ . If L is even, then L is integral and  $L \subset L^*$ .

Let  $g \in O(L)$ . We define the *fixed-point sublattice*  $L^g$  of g and the *coinvariant lattice*  $L_g$  of L associated with g by

(2.1) 
$$L^g = \{ \alpha \in L \mid g\alpha = \alpha \} \text{ and } L_g = \{ \alpha \in L \mid (\alpha | L^g) = 0 \},$$

respectively.

Let L be an even lattice. The discriminant group  $\mathcal{D}(L)$  is defined by  $L^*/L$ , which is a finite abelian group.

Let

(2.2) 
$$q_L: \mathcal{D}(L) \to \mathbb{Q}/\mathbb{Z}, \quad \alpha + L \mapsto \frac{(\alpha | \alpha)}{2} + \mathbb{Z}.$$

Then  $q_L$  defines a quadratic form on  $\mathcal{D}(L)$ . The associated bilinear form on  $\mathcal{D}(L)$  given by  $(\alpha + L|\beta + L) = (\alpha|\beta) + \mathbb{Z}$  is non-degenerate. Hence  $(\mathcal{D}(L), q_L)$  is a non-degenerate quadratic space. An element x of a quadratic space (V, q) is called *singular* if  $x \neq 0$  and q(x) = 0.

**Definition 2.1.** Let p be a prime. An integral lattice L is said to be *p*-elementary if  $pL^* \subset L$ .

We recall a result from [LS24].

**Lemma 2.2.** ([LS24, Lemma 2.1]) Let L be an even lattice with a positive-definite bilinear form  $(\cdot|\cdot)$ . If g is a fixed-point free isometry of L, then

$$((1-g)L^*)^* = (1-g)^{-1}L.$$

The following Lemma will be used later.

**Lemma 2.3.** Let L and g be as in Lemma 2.2. Moreover we assume that  $(1-g)L^* \subset L$ . Then we have

 $nL^* \subset L$ ,

where n is the order of g.

Proof. By the assumption  $(1-g)L^* \subset L$ , it suffices to prove that  $nL^* \subset (1-g)L^*$ . This condition is equivalent to  $(1-g)^{-1}L \subset (1/n)L$ . Since g is fixed-point free, we see that 1-g is non-singular. Moreover, since  $(1-g)(1+g+\cdots+g^{n-1})=0$ , we have  $1+g+\cdots+g^{n-1}=0$ . Since  $(1-g)\sum_{k=1}^{n-1}kg^k = -n$  id, we have  $(1-g)^{-1} = -1/n\sum_{k=1}^{n-1}kg^k$ . Thus we see that  $n(1-g)^{-1}L = n(-1/n\sum_{k=1}^{n-1}kg^k)L \subset L$ , which implies that  $(1-g)^{-1}L \subset (1/n)L$ .

2.2. Notations of Groups. In this subsection, we describe groups which appear in this paper. Let G be a group. For a subgroup H of G, the symbols  $N_G(H)$  and  $C_G(H)$  denote the normalizer of H in G and the centralizer of H in G, respectively. Let n denote the cyclic group with order n. The symbol  $D_n$  denotes the dihedral group with order n. Notations of extraspecial group follow [ATLAS].

According to [Gr02], we explain groups related to orthogonal groups over finite field. Let F be a finite field with  $\operatorname{Char}(F) \neq 2$  and let V be a non-degenerate quadratic space over F. A subspace U of V is called a *totally isotropic subspace* of V if the quadratic form vanishes on U. The dimension of a maximal totally isotropic subspace of V is called the *Witt index* of V, which is denoted by m(V). Then V is expressed as a form  $V = U \oplus W$ , where U is a certain 2m(V)-dimensional subspace of V called hyperbolic and W is a subspace which contains no singular vector. Let  $\dim(V)$  be even. If the above Wvanishes, then the quadratic space V is called (+)-type. Otherwise, V is called (-)-type.

We assume that the Witt index of V is greater than or equal to one. Denote the orthogonal group of the quadratic space V by GO(V). We use  $\Omega(V)$  to denote the commutator subgroup of GO(V) except for the case when  $\dim(V) = 4$  and V is of (+)-type. Set #F = q. If the dimension of V is odd, GO(V) and  $\Omega(V)$  are also written by  $GO_{2m+1}(q)$  and  $\Omega_{2m+1}(q)$ , respectively. Similarly, if the dimension of V is even, then GO(V) and  $\Omega(V)$  are written by  $GO_{2m}^{\epsilon}(q)$  and  $\Omega_{2m}^{\epsilon}(q)$ , where  $\epsilon$  is defined as follows:

$$\epsilon = \begin{cases} + & \text{if } V \text{ is of } (+)\text{-type} \\ - & \text{if } V \text{ is of } (-)\text{-type.} \end{cases}$$

2.3. VOAs, modules and automorphisms. A vertex operator algebra (VOA)  $(V, Y, \mathbb{1}, \omega)$ is a  $\mathbb{Z}$ -graded vector space  $V = \bigoplus_{i \in \mathbb{Z}} V_i$  over the complex field  $\mathbb{C}$  equipped with a linear map

$$Y(a, z) = \sum_{i \in \mathbb{Z}} a_i z^{-i-1} \in (\text{End } V)[[z, z^{-1}]], \quad a \in V,$$

the vacuum vector  $1 \in V_0$  and the conformal vector  $\omega \in V_2$  satisfying some axioms ([FLM88]).

Let V be a VOA. A linear automorphism  $\tau$  of V is called an *automorphism* of V if the linear automorphism satisfies the following:

(1) 
$$\tau \omega = \omega;$$
  
(2)  $\tau Y(v, z) = Y(\tau v, z)\tau$  for all  $v \in V.$ 

The symbol Aut (V) denotes the group of all automorphisms of V. For an automorphism  $\tau$  of V, let  $V^{\tau} = \{v \in V \mid \tau v = v\}$ , which is called the *orbifold VOA*. Then  $V^{\tau}$  is a subVOA of V.

For a VOA V, a V-module  $(M, Y_M)$  is a  $\mathbb{C}$ -graded vector space  $M = \bigoplus_{i \in \mathbb{C}} M_i$  equipped with a linear map

$$Y_M(a,z) = \sum_{i \in \mathbb{Z}} a_i z^{-i-1} \in (\text{End } M)[[z, z^{-1}]], \quad a \in V$$

satisfying some conditions ([FHL93]). We often denote a V-module  $(M, Y_M)$  by M.

Let V be a VOA and let M be an irreducible V-module. Then there exists  $\varepsilon(M) \in \mathbb{C}$ such that  $M = \bigoplus_{m \in \mathbb{Z}_{\geq 0}} M_{\varepsilon(M)+m}$  and  $M_{\varepsilon(M)}$  is not trivial. The number  $\varepsilon(M)$  is called the *conformal weight* of M. For a VOA V, let  $\operatorname{Irr}(V)$  denote the set of all isomorphism classes of irreducible V-modules. We often identify an element in  $\operatorname{Irr}(V)$  with its representative. For irreducible V-modules  $M_1$  and  $M_2$ , under certain assumptions, the fusion rule  $M_1 \boxtimes M_2$ is defined as in [HL95].

**Definition 2.4.** ([DLM00]) Let V be a VOA and  $\tau \in \text{Aut}(V)$ . For a V-module M, the  $\tau$ -conjugate  $(M \circ \tau, Y_{M \circ \tau}(\cdot, z))$  of M is defined as follows:

- (1)  $M \circ \tau = M$  as a vector space;
- (2)  $Y_{M \circ \tau}(a, z) = Y_M(\tau a, z)$  for any  $a \in V$ .

The  $\tau$ -conjugation defines an action of Aut (V) on Irr(V). The following lemma follows by definition.

**Lemma 2.5.** Let  $M, M^1, M^2$  be V-modules and let  $\tau \in Aut(V)$ .

- (1) If M is irreducible, then so is  $M \circ \tau$ . In addition, both M and  $M \circ \tau$  have the same conformal weight.
- (2) Assume that the fusion product is defined on V-modules. Then  $(M^1 \circ \tau) \boxtimes (M^2 \circ \tau) \cong (M^1 \boxtimes M^2) \circ \tau$ . Hence  $\tau$ -conjugation preserves the fusion product.

2.4. Lattice VOAs and their automorphism groups. In this section, we review some facts about lattice VOAs and their automorphism groups from [FLM88, DN99, LY14].

We consider the central extension  $\hat{L}$  of L with the commutator map defined by  $(\cdot|\cdot) \mod 2$ . Let Aut  $(\hat{L})$  be the automorphism group of  $\hat{L}$ . For  $\varphi \in \text{Aut}(\hat{L})$ , we define the element  $\bar{\varphi} \in \text{Aut}(L)$  by  $\bar{\varphi}(\alpha) = \overline{\varphi(e^{\alpha})}$  for  $\alpha \in L$ . We denote  $\{\varphi \in \text{Aut}(\hat{L}) \mid \bar{\varphi} \in O(L)\}$  by  $O(\hat{L})$ 

Let L be a positive-definite even lattice with a positive-definite symmetric bilinear form  $(\cdot|\cdot)$ . Let M(1) be the Heisenberg VOA associated with  $\mathfrak{h} = \mathbb{C} \otimes_{\mathbb{Z}} L$ . Let  $\mathbb{C}\{L\} = \bigoplus_{\alpha \in L} \mathbb{C}e^{\alpha}$  be the twisted group algebra such that  $e^{\alpha}e^{\beta} = (-1)^{(\alpha|\beta)}e^{\beta}e^{\alpha}$  for  $\alpha, \beta \in L$ . The *lattice VOA*  $V_L$  associated with L is defined by  $M(1) \otimes \mathbb{C}\{L\}$  ([FLM88]).

For  $x \in \mathfrak{h}$ , set  $\sigma_x = \exp(-2\pi\sqrt{-1}x_{(0)}) \in \operatorname{Aut}(V_L)$  and let  $N(V_L) = \langle \exp(a_{(0)}) \mid a \in (V_L)_1 \rangle$ . Then  $\sigma_x$  is in  $N(V_L)$  for  $x \in \mathfrak{h}$ . We remark that  $\operatorname{Aut}(V_L) = N(V_L) O(\hat{L})$  ([DN99]). An element  $\phi \in O(\hat{L})$  is called a *standard lift* of  $g \in O(L)$  if  $\bar{\phi} = g$  and  $\phi(e^{\alpha}) = e^{\alpha}$  for  $\alpha \in L^g$ . The following lemma can be found in [EMS20, LS20].

**Lemma 2.6.** Let  $g \in O(L)$  and let  $\hat{g} \in O(\hat{L})$  be a standard lift of g.

(1) Any standard lift of g is conjugate to  $\hat{g}$  by an element in Aut  $(V_L)$ .

(2) If g is fixed-point free or has odd order, then  $|\hat{g}| = |g|$ .

We have exact sequences, which will be used later.

**Theorem 2.7.** ([LY14, Theorem 5.15]) Let L be a positive-definite rootless even lattice. Let g be a fixed-point free isometry of L of prime order p and  $\hat{g}$  a lift of g in  $O(\hat{L})$ . Then we have the exact sequences

$$1 \longrightarrow \operatorname{Hom}(L/(1-g)L, \mathbb{Z}_p) \longrightarrow N_{\operatorname{Aut}(V_L)}(\langle \hat{g} \rangle) \xrightarrow{\varphi} N_{O(L)}(\langle g \rangle) \longrightarrow 1;$$
$$1 \longrightarrow \operatorname{Hom}(L/(1-g)L, \mathbb{Z}_p) \longrightarrow C_{\operatorname{Aut}(V_L)}(\hat{g}) \longrightarrow C_{O(L)}(g) \longrightarrow 1.$$

# 3. Irreducible $V_L^{\hat{g}}$ -modules and their conjugates

Let L be a positive-definite even lattice and let g be a fixed-point free isometry of L of prime order p. Let  $\hat{g} \in O(\hat{L})$  be a standard lift of g. By Lemma 2.6,  $\hat{g}$  also has order p.

In [CM, Mi15], it was proved that the orbifold VOA  $V_L^{\hat{g}}$  is rational,  $C_2$ -cofinite, selfdual and of CFT-type. Under these conditions, the fusion product can be defined on  $V_L^{\hat{g}}$ -modules.

By [DRX17], any irreducible  $V_L^{\hat{g}}$ -module is a submodule of an irreducible  $\hat{g}^s$ -twisted  $V_L$ -module for some  $0 \leq s \leq p-1$ . Regarding definition of twisted module, refer to [DRX17].

**Definition 3.1.** ([LS24, Definition 3.1]) An irreducible  $V_L^{\hat{g}}$ -module is said to be of  $\hat{g}^s$ -type if it is a  $V_L^{\hat{g}}$ -submodule of an irreducible  $\hat{g}^s$ -twisted  $V_L$ -module. In addition, it is said to be of *untwisted type* (resp. *twisted type*) if it is of  $\hat{g}^0$ -type (resp. of  $\hat{g}^s$ -type for some  $1 \leq s \leq p-1$ ).

From here, throughout this section, we assume

$$(3.1) (1-g)L^* \subset L$$

3.1. Irreducible  $V_L^{\hat{g}}$ -modules of untwisted type. In this subsection, we discuss the irreducible  $V_L^{\hat{g}}$ -modules of untwisted type.

Let  $\lambda + L \in \mathcal{D}(L)$  and  $V_{\lambda+L} = M(1) \otimes \operatorname{Span}_{\mathbb{C}} \{ e^{\alpha} \mid \alpha \in \lambda + L \}$ . Then  $V_{\lambda+L}$  has an irreducible  $V_L$ -module structure ([FLM88]). Since  $(1-g)L^* \subset L$ , we see that  $g(\lambda) + L =$ 

 $\lambda + L$ . This implies that  $V_{\lambda+L}$  is  $\hat{g}$ -invariant. Let  $\hat{g}_{\lambda+L}$  be the  $\hat{g}$ -module isomorphism of  $V_{\lambda+L}$ . For  $0 \leq j \leq p-1$ , set

(3.2) 
$$V_{\lambda+L}(j) = \{ v \in V_{\lambda+L} \mid \hat{g}_{\lambda+L}(v) = \exp(2\pi\sqrt{-1}j/p)v \}.$$

We recall some lemmas, which will be used later.

**Lemma 3.2.** ([BLS24, Lemma 3.1]) Let  $h \in N_{O(L)}(g)$  and let  $\hat{h} \in N_{Aut(V_L)}(\hat{g})$  be a lift of h. Let  $\lambda + L \in \mathcal{D}(L)$ . As sets of isomorphism classes of irreducible  $V_L^{\hat{g}}$ -modules, we have

 $\{V_{\lambda+L}(j) \circ \hat{h} \mid 0 \le j \le p-1\} = \{V_{h^{-1}(\lambda)+L}(j) \mid 0 \le j \le p-1\}.$ 

**Lemma 3.3.** ([BLS24, Lemma 3.2]) Let  $\alpha, \lambda \in L^*$  and  $0 \le i, j \le p-1$ . If  $(\alpha|\lambda) \in j/p+\mathbb{Z}$ , then as  $V_L^{\hat{g}}$ -modules,

$$V_{\lambda+L}(i) \circ \sigma_{(1-g)^{-1}\alpha} \cong V_{\lambda+L}(i-j).$$

The following proposition describes the stabilizer of  $V_L(1)$  and the stabilizer of  $\{V_L(i) \mid 0 \le i \le p-1\}$ .

**Proposition 3.4.** ([Sh04, Theorem 3.3])

(1) The stabilizer of  $V_L(1)$  in Aut  $(V_L^{\hat{g}})$  is isomorphic to  $C_{\text{Aut}(V_L)}(\hat{g})/\langle \hat{g} \rangle$ .

(2) The stabilizer of  $\{V_L(i) \mid 0 \le i \le p-1\}$  in Aut  $(V_L^{\hat{g}})$  is isomorphic to  $N_{\text{Aut}(V_L)}(\langle \hat{g} \rangle)/\langle \hat{g} \rangle$ .

3.2. Irreducible  $V_L^{\hat{g}}$ -modules of twisted type. In this subsection, we discuss the irreducible  $V_L^{\hat{g}}$ -modules of twisted type. We use the descriptions in [ALY18, Sections 3.2 and 3.3]. Let  $1 \leq s \leq p-1$ . The irreducible  $\hat{g}^s$ -twisted module  $V_L^T[\hat{g}^s]$  is given by

$$V_L^T[\hat{g}^s] = M(1)[g^s] \otimes T,$$

where  $M(1)[g^s]$  is the  $g^s$ -twisted free bosonic space and T is an irreducible module for a certain  $\hat{g}^s$ -twisted central extension of L.

3.3. Irreducible  $V_L^{\hat{g}}$ -modules and the quadratic form. In this subsection, we summarize the classification of irreducible  $V_L^{\hat{g}}$ -modules and review the quadratic form on  $\operatorname{Irr}(V_L^{\hat{g}})$ . For the irreducible  $V_L^{\hat{g}}$ -modules, we adopt the notation in [La20].

The following theorem describes the structure of  $\operatorname{Irr}(V_L^{\hat{g}})$ .

**Theorem 3.5** ([La20, Theorem 5.3]). Let L be a positive-definite even lattice and let g be a fixed-point free isometry of L of odd prime order p. Assume that  $(1-g)L^* \subset L$ . Then

 $\operatorname{Irr}(V_L^{\hat{g}}) = \{ V_{\lambda+L}[\hat{g}^i](j) \mid \lambda + L \in \mathcal{D}(L), \ 0 \le i, j \le p-1 \},\$ 

where  $V_{\lambda+L}[\hat{g}^i](j)$  is an irreducible  $V_L^{\hat{g}}$ -module and we identify  $V_{\lambda+L}(j)$  with  $V_{\lambda+L}[\hat{g}^0](j)$ . In particular,  $V_L^{\hat{g}}$  has exactly  $|\mathcal{D}(L)|p^2$  inequivalent irreducible modules. Furthermore, if the conformal weight of  $V_L[\hat{g}]$  is in  $(1/p)\mathbb{Z}$ , then  $\operatorname{Irr}(V_L^{\hat{g}}) \cong \mathcal{D}(L) \times (\mathbb{Z}/p\mathbb{Z})^2$  as abelian groups.

The explicit fusion products are given as follows:

(3.3) 
$$V_{\lambda_1+L}[\hat{g}^{i_1}](j_1) \boxtimes V_{\lambda_2+L}[\hat{g}^{i_2}](j_2) = V_{\lambda_1+\lambda_2+L}[\hat{g}^{i_1+i_2}](j_1+j_2).$$

By the theorem above,  $\operatorname{Irr}(V_L^{\hat{g}})$  forms a finite abelian group under the fusion product. Let

(3.4) 
$$q: \operatorname{Irr}(V_L^{\hat{g}}) \to \mathbb{Q}/\mathbb{Z}, \quad M \mapsto \varepsilon(M) \pmod{\mathbb{Z}}.$$

The conformal weight of  $V_L^{\hat{g}}$  is the following.

$$\varepsilon(V_{\lambda+L}[\hat{g}^i](j)) \equiv \frac{ij}{p} + \frac{(\lambda|\lambda)}{2} \pmod{\mathbb{Z}}.$$

Then the form  $\langle \mid \rangle : \operatorname{Irr}(V_L^{\hat{g}}) \times \operatorname{Irr}(V_L^{\hat{g}}) \to \mathbb{Q}/\mathbb{Z}$  associated with q is defined by

(3.5) 
$$\langle M^1 | M^2 \rangle = q(M^1 \boxtimes M^2) - q(M^1) - q(M^2) \pmod{\mathbb{Z}}.$$

The following theorem describes the structure of the quadratic space.

**Theorem 3.6.** ([EMS20, Theorem 3.4]) ( $\operatorname{Irr}(V_L^{\hat{g}}), q$ ) is a non-degenerate quadratic space, that is, the form  $\langle | \rangle$  is non-degenerate and bilinear.

To examine certain orbits, we will use the following proposition later.

**Proposition 3.7.** ([LS24, Lemma 3.5]) Let  $\alpha \in L^*$  and  $1 \leq i \leq p-1$ . For  $0 \leq j \leq p-1$ ,

$$V_L[\hat{g}^i](j) \circ \sigma_{(1-g^i)^{-1}\alpha} \cong V_{\alpha+L}[\hat{g}^i](j'),$$

where j' is determined by  $ij \equiv p(\alpha|\alpha)/2 + ij' \pmod{p}$ . In particular, all irreducible  $V_L^{\hat{g}}$ -modules of  $\hat{g}^i$ -type with the same conformal weight are conjugate under the action of Aut  $(V_L^{\hat{g}})$ .

# 4. Construction B associated with codes over $\mathbb{Z}_p$

In this section, according to [LS24], we review Construction B associated with codes over  $\mathbb{Z}_p$ .

Let  $R_i$   $(1 \le i \le t)$  be a copy of the root lattice of type  $A_{p-1}$ . The following is a realization of the root lattice of type  $A_{p-1}$ :

$$\left\{ (a_1, a_2, \dots, a_p) \in \mathbb{Z}^p \mid \sum_{i=1}^p a_i = 0 \right\}.$$

For  $1 \leq k \leq p$ , we define  $v_k \in \mathbb{Z}^p$  by the vector

$$(0,\ldots,0,\overset{k}{1},0,\ldots,0).$$

Set  $\alpha_k = v_k - v_{k+1}$  for  $1 \le k \le p-1$ . Then  $\Delta = \{\alpha_k \mid 1 \le k \le p-1\}$  is a base of the root system of  $A_{p-1}$ . Let  $\Delta_+$  be the set of all positive roots with respect to the base  $\Delta$ . The vector  $1/2 \sum_{\alpha \in \Delta_+} \alpha$  is called the *Weyl vector* with respect to the base  $\Delta$ . The *Coxeter element* with respect to the above base  $\Delta$  is defined by the transformation on  $\mathbb{Z}^p$ :

$$(x_1, x_2, \dots, x_p) \mapsto (x_p, x_1, x_2, \dots, x_{p-1}).$$

Set  $R = R_1 \perp \cdots \perp R_t$ . Then  $\mathcal{D}(R) \cong \mathbb{Z}_p^t$ . For each  $1 \leq i \leq t$ , fix the above base  $\Delta_i$  of the root system  $R_i(2)$  of type  $A_{p-1}$ , and set  $\Delta = \bigcup_{i=1}^t \Delta_i$ .

Let

$$\nu: R^* \to R^*/R = \mathcal{D}(R) \cong \mathbb{Z}_p^t$$

be the canonical surjective map. For a subgroup C of  $\mathbb{Z}_p^t$ , let  $L_A(C)$  denote the lattice defined by

(4.1) 
$$L_A(C) = \nu^{-1}(C) = \{ \alpha \in R^* \mid \nu(\alpha) \in C \};$$

we call  $L_A(C)$  the lattice constructed by Construction A from C.

Let  $g_{\Delta_i} \in O(R_i)$  be the Coxeter element defined by the base  $\Delta_i$ . For  $e = (e_i) \in \mathbb{Z}_p^t$ , set

(4.2) 
$$g_{\Delta,e} = ((g_{\Delta_1})^{e_1}, \dots, (g_{\Delta_t})^{e_t}) \in O(L_A(C)).$$

If  $gcd(e_i, p) = 1$  for all  $1 \le i \le t$ , then the map  $g_{\Delta,e}$  is fixed-point free.

Set

(4.3) 
$$\chi_{\Delta} = \frac{1}{p}(\rho_{\Delta_1}, \dots, \rho_{\Delta_t}) \in \mathbb{Q} \otimes_{\mathbb{Z}} R$$

where  $\rho_{\Delta_i}$  is the Weyl vector of  $R_i$  with respect to  $\Delta_i$ .

Let  $L_B(C)$  be the sublattice of  $L_A(C)$  defined by

(4.4) 
$$L_B(C) = \{ \alpha \in L_A(C) \mid (\alpha | \chi_{\Delta}) \in \mathbb{Z} \};$$

we call  $L_B(C)$  the lattice constructed by Construction B from C.

Let  $\lambda_0 = 0$  and define  $\lambda_l$  for  $1 \le l \le p - 1$  as follows:

$$\lambda_l = \frac{1}{p} \left( (p-l) \sum_{k=1}^l v_k - l \sum_{k=l+1}^p v_k \right).$$

For  $x = (x_i) \in \mathbb{Z}_p^t$ , denote

(4.5) 
$$\lambda_x = (\lambda_{x_1}^1, \dots, \lambda_{x_t}^t) \in R_1^* \perp \dots \perp R_t^* = R^*,$$

where  $x_i$  is regarded as an element of  $\{0, \ldots, p-1\}$ .

The following results were proved in [LS24].

**Lemma 4.1.** ([LS24, Lemma 4.10])  $g_{\Delta,e} \in O(L_B(C))$  if and only if  $\lambda_e \in L_A(C)^*$ .

**Lemma 4.2.** ([LS24, Lemma 4.13]) Let C be a self-orthogonal code over  $\mathbb{Z}_p$ . Then  $L_A(C)^* = L_A(C^{\perp})$ .

5. The structure of the kernel associated with the action of Aut  $(V_L^{\hat{g}})$ 

Let L be a positive-definite even lattice and let g be a fixed-point free isometry of L of prime order p. In this section, we consider the kernel of the action of Aut  $(V_L^{\hat{g}})$  on  $\operatorname{Irr}(V_L^{\hat{g}})$ .

We review the action of  $\operatorname{Aut}(V_L^{\hat{g}})$  on  $\operatorname{Irr}(V_L^{\hat{g}})$ . By Subsection 2.3,  $\operatorname{Aut}(V_L^{\hat{g}})$  acts on  $\operatorname{Irr}(V_L^{\hat{g}})$  by the conjugate action. Moreover, the action preserves the quadratic form of  $\operatorname{Irr}(V_L^{\hat{g}})$  as described in Theorem 3.6. Hence we can consider the group homomorphism

(5.1) 
$$\mu \colon \operatorname{Aut} (V_L^{\hat{g}}) \longrightarrow O(\operatorname{Irr}(V_L^{\hat{g}}), q).$$

We determine the structure of the kernel of  $\mu$ . The following is a generalization of [Sh04, Lemma 3.7].

**Theorem 5.1.** Let L be a positive-definite rootless even lattice and let g be a fixed-point free isometry of L of prime order p. If  $(1-g)L^* \subset L$ , then we have the following exact sequence

$$1 \longrightarrow A \longrightarrow \operatorname{Ker} \mu \stackrel{\tilde{\varphi}}{\longrightarrow} B \longrightarrow 1,$$

where  $A = \text{Hom}(L/(1-g)L^*, \mathbb{Z}_p)$  and  $B = \{h \in C_{O(L)}(g) \mid h = 1 \text{ on } L^*/L\}/\langle g \rangle$ .

*Proof.* By Theorem 2.7, we have an exact sequence

(5.2) 
$$1 \longrightarrow \operatorname{Hom}(L/(1-g)L, \mathbb{Z}_p) \longrightarrow C_{\operatorname{Aut}(V_L)}(\hat{g})/\langle \hat{g} \rangle \xrightarrow{\tilde{\varphi}} C_{O(L)}(g)/\langle g \rangle \longrightarrow 1.$$

We first determine  $\operatorname{Hom}(L/(1-g)L, \mathbb{Z}_p) \cap \operatorname{Ker} \mu$ . By Lemma 3.3, if  $(\alpha|\lambda) \in j/p + \mathbb{Z}$  for  $\alpha, \lambda \in L^*$ , then as  $V_L^{\hat{g}}$ -modules,

$$V_{\lambda+L}(i) \circ \sigma_{(1-g)^{-1}\alpha} \cong V_{\lambda+L}(i-j).$$

By Lemma 2.3, L is p-elementary. Hence, for any  $\alpha, \lambda \in L^*$ , we have

$$(\alpha|\lambda) \in \{i/p + \mathbb{Z} \mid 0 \le i \le p - 1\}.$$

Hence, if  $\alpha$  is an element of  $\operatorname{Hom}(L/(1-g)L, \mathbb{Z}_p) \cap \operatorname{Ker} \mu$ , then we have  $\alpha \in (L^*)^* = L$ . Thus we see that  $\operatorname{Hom}(L/(1-g)L, \mathbb{Z}_p) \cap \operatorname{Ker} \mu \subset \{\sigma_\alpha \mid \alpha \in (1-g)^{-1}L/L^*\}$ . Conversely, by the proof of [LY14, Theorem 5.15], for  $\alpha \in (1-g)^{-1}L/L^*$ ,  $\sigma_\alpha$  corresponds to the map defined by  $\beta \mapsto (p\alpha|\beta) \mod p$  for  $\beta \in L$ . Since  $p\alpha \in p(1-g)^{-1}L \subset L$ , for any twisted module  $V_{\lambda+L}[\hat{g}^i](j)$ , we have  $V_{\lambda+L}[\hat{g}^i](j) \circ \sigma_{\alpha} \cong V_{\alpha+\lambda+L}[\hat{g}^i](j) \cong V_{\lambda+L}[\hat{g}^i](j)$ . Hence,  $\sigma_{\alpha}$  fixes all irreducible  $V_L^{\hat{g}}$ -modules of twisted type. Moreover, by Lemma 3.3, we see that  $\sigma_{\alpha}$  fixes all irreducible  $V_L^{\hat{g}}$ -modules of untwisted type. Hence we have  $\operatorname{Hom}(L/(1-g)L, \mathbb{Z}_p) \cap$ Ker  $\mu = \{\sigma_{\alpha} \mid \alpha \in (1-g)^{-1}L/L^*\}$ . Note that  $\operatorname{Hom}(L/(1-g)L^*, \mathbb{Z}_p) \cong \{\sigma_{\alpha} \mid \alpha \in (1-g)^{-1}L/L^*\}$ .

Next we prove that the map  $\tilde{\varphi}$  from Ker  $\mu$  to B is surjective. By Lemma 3.2, for any  $\lambda \in L^*$  and  $0 \leq j \leq p-1$ , there exists  $0 \leq j' \leq p-1$  such that

$$V_{\lambda+L}(j) \circ \hat{h} \cong V_{h^{-1}(\lambda)+L}(j'),$$

where  $\hat{h} \in C_{\operatorname{Aut}(V_L)}(\hat{g})/\langle \hat{g} \rangle$  and  $\tilde{\varphi}(\hat{h}) = h$ . Hence, if  $\tau \in \operatorname{Ker} \mu$ , then for any  $\lambda \in \mathcal{D}(L)$ , we have  $\varphi(\tau)^{-1}(\lambda) - \lambda \in L$ , which means that  $\varphi(\tau)$  is 1 on  $\mathcal{D}(L)$ . Thus we have  $\tilde{\varphi}(\operatorname{Ker} \mu) \subset B$ .

Conversely, let  $h \in B$ . By Theorem 2.7, we obtain an element  $\hat{h}$  of  $C_{\text{Aut}(V_L^{\hat{g}})}(\hat{g})$  such that  $\varphi(\hat{h}) = h$ . By Lemma 3.2, for any  $\lambda + L \in \mathcal{D}(L)$ , there exists  $0 \leq j \leq p - 1$  such that

$$V_{\lambda+L}(0) \circ \hat{h} \cong V_{\lambda+L}(j).$$

Denote the above j for  $\lambda + L \in \mathcal{D}(L)$  by  $j_{\lambda+L}$ . Then we can define a map f from  $\mathcal{D}(L)$  to  $\mathbb{Z}/p\mathbb{Z}$  defined by  $\lambda + L \mapsto j_{\lambda+L}$ . This map f is a group homomorphism. In fact, by Lemma 2.5 (2), we have

$$(V_{\lambda_1+L}(0) \boxtimes V_{\lambda_2+L}(0)) \circ \hat{h} \cong (V_{\lambda_1+L}(0) \circ \hat{h}) \boxtimes (V_{\lambda_2+L}(0) \circ \hat{h}) \text{ for } \lambda_1, \lambda_2 \in \mathcal{D}(L).$$

By (3.3), we have  $j_{\lambda_1+L} + j_{\lambda_2+L} = j_{(\lambda_1+\lambda_2)+L}$ , which means that the map f is a group homomorphism. By Lemma 2.3, L is *p*-elementary. Hence, we can consider an injective group homomorphism

$$\mathcal{D}(L) \longrightarrow \operatorname{Hom}(L^*/L, \mathbb{Z}_p), \ \alpha \mapsto p(\alpha|\cdot) \mod p.$$

Moreover, since  $\mathcal{D}(L)$  is a finite dimensional vector space over  $\mathbb{F}_p$ , the above group homomorphism is bijective. Thus there exists an element  $\beta$  of  $\mathcal{D}(L)$  such that  $f = -p(\beta|\cdot) \mod p$ . Define  $\hat{h}_0 = \sigma_{(1-g)^{-1}\beta}\hat{h}$ . By Lemma 3.2, for any  $\lambda + L \in \mathcal{D}(L)$ , we have

$$V_{\lambda+L}(0) \circ \hat{h}_0 \cong V_{\lambda+L}(0).$$

The orthogonal complement of the subspace of  $\{V_{\lambda+L}(0) \mid \lambda + L \in \mathcal{D}(L)\}$  in  $\operatorname{Irr}(V_L^{\hat{g}})$  with respect to the bilinear form in (3.5) is  $\{V_L[\hat{g}^i](j) \mid 0 \leq i, j \leq p-1\}$ , which is preserved by the action of  $\hat{h}_0$ . Since  $\hat{h}$  and  $\hat{g}$  are commutative,  $\hat{h}_0$ -conjugates of  $\hat{g}$ -modules are also  $\hat{g}$ -modules. Hence, by Proposition 3.7, we have

$$V_L[\hat{g}](0) \circ \hat{h}_0 \cong V_L[\hat{g}](0).$$

By Theorem 3.5,  $\operatorname{Irr}(V_L^{\hat{g}})$  is generated by  $\{V_L(j), V_{\lambda+L}(0), V_L[\hat{g}](0) \mid 0 \leq j \leq p-1, \lambda \in \mathcal{D}(L)\}$ . Since the above generators are preserved by the action of  $\hat{h}_0$ , by Lemma 2.5 (2), all elements of  $\operatorname{Irr}(V_L^{\hat{g}})$  are preserved by the action of  $\hat{h}_0$ . This means that  $\hat{h}_0 \in \operatorname{Ker} \mu$ , as desired.

By using Theorem 5.1, we have the following corollary.

**Corollary 5.2.** Let L be a positive-definite rootless even lattice and let g be a fixed-point free isometry of L of prime order p. Let  $\mu$  be as in (5.1). If  $(1 - g)L^* \subset L$ , then the following sequence is exact

(5.3) 
$$1 \longrightarrow H_1/H_2 \longrightarrow \operatorname{Stab}_{\operatorname{Im}\mu}(V_L(1)) \xrightarrow{\tilde{\varphi}'} K_1/K_2 \longrightarrow 1,$$

where  $H_1 = \text{Hom}(L/(1-g)L, \mathbb{Z}_p), H_2 = \text{Hom}(L/(1-g)L^*, \mathbb{Z}_p), K_1 = C_{O(L)}(g)/\langle g \rangle$  and  $K_2 = \{h \in C_{O(L)}(g) \mid h = 1 \text{ on } L^*/L\}/\langle g \rangle.$ 

Proof. By Proposition 3.4, we can identify  $\operatorname{Stab}_{\operatorname{Aut}(V_L^{\hat{g}})}(V_L(1))$  with  $C_{\operatorname{Aut}(V_L)}(\hat{g})/\langle \hat{g} \rangle$ . Let  $\pi_1$  be the natural map from  $\operatorname{Stab}_{\operatorname{Aut}(V_L^{\hat{g}})}(V_L(1))$  to  $\operatorname{Stab}_{\operatorname{Aut}(V_L^{\hat{g}})}(V_L(1))/\operatorname{Ker}\mu$  and let  $\pi_2$  be the natural map from  $K_1$  to  $K_1/K_2$ . Let i be the inclusion in (5.2) and let  $\tilde{\varphi}$  be the surjective map in (5.2). By Theorem 5.1, we see that  $\operatorname{Ker}(\pi_1 \circ i) = H_2$ , which means that we have the injection map from  $H_1/H_2$  to  $\operatorname{Stab}_{\operatorname{Aut}(V_L^{\hat{g}})}(V_L(1))/\operatorname{Ker}\mu$ . Moreover, we see that  $\operatorname{Ker}(\pi_2 \circ \tilde{\varphi}) \supset \operatorname{Ker}\mu$ , which implies that we have the surjective map  $\tilde{\varphi}'$  from  $\operatorname{Stab}_{\operatorname{Aut}(V_T^{\hat{g}})}(V_L(1))/\operatorname{Ker}\mu$  to  $K_1/K_2$ . Hence we have the following sequence

(5.4) 
$$1 \longrightarrow H_1/H_2 \longrightarrow \operatorname{Stab}_{\operatorname{Aut}(V_L^{\hat{g}})}(V_L(1))/\operatorname{Ker} \mu \xrightarrow{\tilde{\varphi}'} K_1/K_2 \longrightarrow 1.$$

Note that the image of  $H_1/H_2$  is included in Ker  $\tilde{\varphi}'$ .

Finally, we verify that  $\operatorname{Ker} \tilde{\varphi}'$  is equal to the image of  $H_1/H_2$ . By isomorphism theorems, we have  $\# \operatorname{Ker} \tilde{\varphi}' = \# \operatorname{Stab}_{\operatorname{Aut}(V_L^{\hat{g}})}(V_L(1))/\# \operatorname{Ker} \mu \cdot \# K_2/\# K_1 = (\# H_1 \# K_1/\# H_2 \# K_2) \cdot \# K_2/\# K_1 = \# H_1/\# H_2$ . Since the image of  $H_1/H_2$  is included in  $\operatorname{Ker} \tilde{\varphi}'$ , we have  $\operatorname{Ker} \tilde{\varphi}'$  is equal to the image of  $H_1/H_2$ . Since  $\operatorname{Stab}_{\operatorname{Aut}(V_L^{\hat{g}})}(V_L(1))/\operatorname{Ker} \mu \cong \operatorname{Stab}_{\operatorname{Im} \mu}(V_L(1))$  as groups, we have the desired result.  $\Box$ 

6. STRUCTURE OF Aut  $(V_L^{\hat{g}})$  in the case  $L = \Lambda_{3C}, \Lambda_{5C}, \Lambda_{11A}$  or  $\Lambda_{23A}$ .

Thoughout this section, we consider the structure of automorphism groups of orbifold vertex operator algebras in the cases  $L = \Lambda_{3C}, \Lambda_{5C}, \Lambda_{11A}$  or  $\Lambda_{23A}$ . To compute the automorphism groups, we use MAGMA in this section ([BCP97]).

6.1. The case  $L = \Lambda_{3C}$ . Let C be the code over  $\mathbb{Z}_3$  with the generator matrix

This code was given in [LS24, Theorem 7.16].

Remark 6.1. By [LS24, Table 1 in Subsection 4.5], we see that  $\Lambda_{3C} \cong L_B(C)$  as lattices and  $\mathcal{D}(L) \cong 3^5$ . Hence,  $\Lambda_{3C}$  is realized by Construction B in Section 4.

For e = (1, 1, 1, 1, 1, 1, 1, 1, 1), we consider the fixed-point free isometry  $g = g_{\Delta,e}$  of Lin Section 4. Note that g is a fixed-point free isometry of  $L_B(C)$  by Lemma 4.1 and Lemma 4.2.

Set  $\mathcal{L}_{3C} = \{\lambda + L \in \mathcal{D}(L) \mid q(\lambda + L) = 0\} \setminus \{L\}$ . By using MAGMA, we can count the number of orbits under the action  $C_{O(L)}(g)/\langle g \rangle$ . Hence we have the following proposition.

# **Proposition 6.2.** $C_{O(L)}(g)/\langle g \rangle$ acts transitively on $\mathcal{L}_{3C}$ .

Let  $S_{3C}$  be the set of all singular vectors in  $\operatorname{Irr}(V_L^{\hat{g}})$ . By Theorem 3.5 and Remark 6.1, we have  $\operatorname{Irr}(V_L^{\hat{g}}) \cong 3^7$  as abelian groups. Hence, by [Wi09, (3.27)], we have  $\#S_{3C} = 3^6 - 1$ . Since  $\Lambda_{3C}$  is realized by Construction B and  $C_{O(L)}(g)/\langle g \rangle$  acts transitively on  $\mathcal{L}_{3C}$ , we can imitate the proof of [BLS24, PROPOSITION 6.5]. Hence we obtain the following proposition.

# **Proposition 6.3.** Aut $(V_L^{\hat{g}})$ also acts transitively on $S_{3C}$ .

Next, we consider the structure of Ker  $\mu$  in (5.1). By Theorem 5.1, we examine  $\operatorname{Hom}(L/(1-g)L^*,\mathbb{Z}_3)$  and  $\{h \in C_{O(L)}(g) \mid h = 1 \text{ on } L^*/L\}/\langle g \rangle$ . By using MAGMA, we see that  $\operatorname{Hom}(L/(1-g)L^*,\mathbb{Z}_3) \cong 3^4$ . Define the matrix group  $T_{3C}$  by the following:

$$\left\{ \begin{pmatrix} 1 & a & b & c \\ 0 & f & 0 & d \\ 0 & 0 & f & e \\ 0 & 0 & 0 & 1 \end{pmatrix} \in \operatorname{GL}_4(\mathbb{F}_3) \mid a, b, c, d, e \in \mathbb{F}_3, f \in \mathbb{F}_3 \setminus \{0\} \right\}.$$

By using MAGMA, we see that  $\{h \in C_{O(L)}(g) \mid h = 1 \text{ on } L^*/L\} \cong T_{3C} \cong 3^{1+4}_+ : 2$ and  $\{h \in C_{O(L)}(g) \mid h = 1 \text{ on } L^*/L\}/\langle g \rangle \cong 3^4 : 2$ . Combining the obtained results with Theorem 5.1, we have the following proposition.

**Proposition 6.4.** Hom $(L/(1-g)L^*, \mathbb{Z}_3) \cong 3^4$  and  $\{h \in C_{O(L)}(g) \mid h = 1 \text{ on } L^*/L\}/\langle g \rangle \cong 3^4 : 2$ . Hence, we have Ker  $\mu \cong 3^4 . (3^4 : 2)$  and  $\# \text{Ker } \mu = 2 \cdot 3^8$ .

Next we consider the structure of Im  $\mu$  in (5.1). First we determine the order of Aut  $(V_L^{\hat{g}})$ . By using MAGMA, we have  $\text{Hom}(L/(1-g)L,\mathbb{Z}_3) \cong 3^9$  and  $\#C_{O(L)}(g)/\langle g \rangle = 2^8 \cdot 3^8 \cdot 5$ . Combining this with Theorem 2.7 and Proposition 3.4, we have the following lemma.

**Proposition 6.5.** Hom $(L/(1-g)L, \mathbb{Z}_3) \cong 3^9$  and  $\#C_{O(L)}(g)/\langle g \rangle = 2^8 \cdot 3^8 \cdot 5$ . Hence  $\# \operatorname{Stab}_{\operatorname{Aut}(V_L^{\hat{g}})}(V_L(1)) = 2^8 \cdot 3^{17} \cdot 5$ .

Next, we consider the order of Aut  $(V_L^{\hat{g}})$ .

**Proposition 6.6.** #Aut  $(V_L^{\hat{g}}) = 2^{11} \cdot 3^{17} \cdot 5 \cdot 7 \cdot 13$ . Hence, by Proposition 6.4, we have  $\# \text{Im } \mu = 2^{10} \cdot 3^9 \cdot 5 \cdot 7 \cdot 13$ .

*Proof.* By Theorem 3.5 and Remark 6.1, we have  $Irr(V_L^{\hat{g}}) \cong 3^7$  as abelian groups. Combining this with Proposition 6.3 and Proposition 6.5, we have

$$#\operatorname{Aut}(V_L^{\hat{g}}) = #S_{3C} # \operatorname{Stab}_{\operatorname{Aut}(V_L^{\hat{g}})}(V_L(1)) = (3^6 - 1) \cdot 2^8 \cdot 3^{17} \cdot 5 = 2^{11} \cdot 3^{17} \cdot 5 \cdot 7 \cdot 13.$$

To determine the structure of Im  $\mu$ , we need to consider  $\operatorname{Stab}_{\operatorname{Im}\mu}(V_L(1))$ . By using MAGMA, we see that  $C_{O(L)}(g)/\{h \in C_{O(L)}(g) \mid h = 1 \text{ on } L^*/L\}$  is isomorphic to a subgroup of  $\operatorname{GO}_5(3)$  of index 2 containing -1, where -1 is the diagonal matrix whose all diagonal entries are -1. Such a subgroup of  $\operatorname{GO}_5(3)$  is determined uniquely and is precisely the group  $2 \times \Omega_5(3)$ . Hence we have  $C_{O(L)}(g)/\{h \in C_{O(L)}(g) \mid h = 1 \text{ on } L^*/L\} \cong 2 \times \Omega_5(3)$ . By Proposition 6.4 and 6.5, we see that  $\operatorname{Hom}(L/(1-g)L, \mathbb{Z}_3)/\operatorname{Hom}(L/(1-g)L^*, \mathbb{Z}_3) \cong 3^5$ . Combining these with Corollary 5.2, we have the following proposition.

## **Proposition 6.7.** Stab<sub>Im $\mu$ </sub>( $V_L(1)$ ) $\cong$ 3<sup>5</sup>. (2 × $\Omega_5(3)$ ).

Finally, we determine the structure of Im  $\mu$ . Since  $\operatorname{Irr}(V_L^{\hat{g}}) \cong 3^7$  as abelian groups, Im  $\mu$  is a subgroup of  $\operatorname{GO}_7(3)$ . By [Wi09], we can compute the order of  $\operatorname{GO}_7(3)$ . Combining this with Proposition 6.6, we see that  $|\operatorname{GO}_7(3) : \operatorname{Im} \mu| = 2$ . We summarize our results. The group Im  $\mu$  is a subgroup of  $\operatorname{GO}_7(3)$  satisfying the following:

- (1)  $|\text{GO}_7(3) : \text{Im } \mu| = 2;$
- (2) Im  $\mu$  acts transitively on  $S_{3C}$ ;
- (3) The stabilizer of an element in  $S_{3C}$  in Im  $\mu$  is isomorphic to  $3^5 \cdot (2 \times \Omega_5(3))$ .

There exist precisely three subgroups of  $GO_7(3)$  of index 2. These three groups are  $SO_7(3), 2 \times \Omega_7(3)$  and  $\Omega_7(3) \cdot 2$ . By using MAGMA, we see that among the above three groups, there exists only one group which satisfies the condition (2) and (3). The group is  $\Omega_7(3) \cdot 2$ . Combining this with Proposition 6.4, we have the following theorem.

**Theorem 6.8.** Aut  $(V_{\Lambda_{3C}}^{\hat{g}}) \cong (3^4 \cdot (3^4 \cdot 2)) \cdot (\Omega_7(3) \cdot 2).$ 

6.2. The case  $L = \Lambda_{5C}$ . This case is similar to the case  $L = \Lambda_{3C}$ . Let C be the code over  $\mathbb{Z}_5$  with the generator matrix

$$\begin{pmatrix} 1 & 1 & 1 & 1 & 1 \\ 1 & 2 & 4 & 3 & 0 \end{pmatrix}.$$

This code was also given in [LS24, Theorem 7.16].

Remark 6.9. By [LS24, Table 1 in Subsection 4.5], we see that  $\Lambda_{5C} \cong L_B(C)$  as lattices and  $\mathcal{D}(L) \cong 5^3$ . Hence,  $\Lambda_{5C}$  is also realized by Construction B in Section 4.

For e = (1, 1, 1, 1, 1), we consider the fixed-point free isometry  $g = g_{\Delta,e}$  of L in Section 4. Note that g is a fixed-point free isometry of  $L_B(C)$  by Lemma 4.1 and Lemma 4.2.

Set  $\mathcal{L}_{5C} = \{\lambda + L \in \mathcal{D}(L) \mid q(\lambda + L) = 0\} \setminus \{L\}$ . By using MAGMA, we can count the number of orbits under the action  $N_{O(L)}(g)/\langle g \rangle$ . Hence we have the following proposition.

### **Proposition 6.10.** $N_{O(L)}(g)/\langle g \rangle$ acts transitively on $\mathcal{L}_{5C}$ .

Remark 6.11. By using MAGMA, we see that  $C_{O(L)}(g)/\langle g \rangle$  does not act transitively on  $\mathcal{L}_{5C}$ . More precisely, the number of orbits is two.

Let  $S_{5C}$  be the set of all singular vectors in  $\operatorname{Irr}(V_L^{\hat{g}})$ . By Theorem 3.5 and Remark 6.9, we have  $\operatorname{Irr}(V_L^{\hat{g}}) \cong 5^5$  as abelian groups. Hence, by [Wi09, (3.27)], we have  $\#S_{5C} = 5^4 - 1$ . Since  $\Lambda_{5C}$  is also realized by Construction B, we can imitate the proof of [BLS24, PROPOSITION 6.5]. Hence we obtain the following proposition.

**Proposition 6.12.** Aut  $(V_L^{\hat{g}})$  also acts transitively on  $S_{5C}$ .

Next, we consider the structure of Ker  $\mu$  in (5.1). By Theorem 5.1, we examine  $\operatorname{Hom}(L/(1-g)L^*,\mathbb{Z}_5)$  and  $\{h \in C_{O(L)}(g) \mid h = 1 \text{ on } L^*/L\}/\langle g \rangle$ . By using MAGMA, we have  $\operatorname{Hom}(L/(1-g)L^*,\mathbb{Z}_5) \cong 5^2$ . Define the matrix group  $T_{5C}$  by the following:

$$\left\{ \begin{pmatrix} 1 & a & b \\ 0 & d & c \\ 0 & 0 & 1 \end{pmatrix} \in \operatorname{GL}_3(\mathbb{F}_5) \mid a, b, c \in \mathbb{F}_5, d \in \{1, -1\} \right\}.$$

By using MAGMA, we see that  $\{h \in C_{O(L)}(g) \mid h = 1 \text{ on } L^*/L\} \cong T_{5C} \cong 5^{1+2}_+ : 2$  and  $\{h \in C_{O(L)}(g) \mid h = 1 \text{ on } L^*/L\}/\langle g \rangle \cong 5^2 : 2$ . Combining obtained results with Theorem 5.1, we have the following proposition.

**Proposition 6.13.** Hom $(L/(1-g)L^*, \mathbb{Z}_5) \cong 5^2$  and  $\{h \in C_{O(L)}(g) \mid h = 1 \text{ on } L^*/L\}/\langle g \rangle \cong 5^2$ : 2. Hence, we have Ker  $\mu \cong 5^2$ .  $(5^2:2)$  and  $\# \text{Ker } \mu = 2 \cdot 5^4$ .

Next we consider the structure of  $\text{Im}\,\mu$  in (5.1). First we determine the order of Aut  $(V_L^{\hat{g}})$ . By using MAGMA, we have  $\operatorname{Hom}(L/(1-g)L, \mathbb{Z}_5) \cong 5^5$  and  $\#C_{O(L)}(g)/\langle g \rangle =$  $2^4 \cdot 3 \cdot 5^3$ . Combining this with Theorem 2.7 and Proposition 3.4, we have the following lemma.

**Proposition 6.14.** Hom $(L/(1-g)L, \mathbb{Z}_5) \cong 5^5$  and  $\#C_{O(L)}(g)/\langle g \rangle = 2^4 \cdot 3 \cdot 5^3$ . Hence we have  $\# \operatorname{Stab}_{\operatorname{Aut}(V_L^{\hat{g}})}(V_L(1)) = 2^4 \cdot 3 \cdot 5^8.$ 

Imitating the proof of Proposition 6.6, we have the following proposition.

**Proposition 6.15.** #Aut $(V_L^{\hat{g}}) = 2^8 \cdot 3^2 \cdot 5^8 \cdot 13$ . Hence, by Proposition 6.13, we have  $\# \operatorname{Im} \mu = 2^7 \cdot 3^2 \cdot 5^4 \cdot 13.$ 

*Proof.* By Theorem 3.5 and Remark 6.9, we have  $Irr(V_L^{\hat{g}}) \cong 5^5$  as abelian groups. Combining this with Proposition 6.12 and Proposition 6.14, we have

$$#\operatorname{Aut}(V_L^{\hat{g}}) = #S_{5C} # \operatorname{Stab}_{\operatorname{Aut}(V_L^{\hat{g}})}(V_L(1)) = (5^4 - 1) \cdot 2^4 \cdot 3 \cdot 5^8 = 2^8 \cdot 3^2 \cdot 5^8 \cdot 13.$$

By using MAGMA, we see that  $C_{O(L)}(g)/\{h \in C_{O(L)}(g) \mid h = 1 \text{ on } L^*/L\}$  is isomorphic to a subgroup of  $GO_3(5)$  of index 2 containing -1, where -1 is the diagonal matrix whose all diagonal entries are -1. Such a subgroup of  $GO_3(5)$  is determined uniquely and is precisely the group  $2 \times \Omega_3(5)$ . Hence we have  $C_{O(L)}(g)/\{h \in C_{O(L)}(g) \mid h = 1 \text{ on } L^*/L\} \cong$  $2 \times \Omega_3(5)$ . By Proposition 6.13 and 6.14, we see that  $\operatorname{Hom}(L/(1-g)L, \mathbb{Z}_5)/\operatorname{Hom}(L/(1-g)L, \mathbb{Z}_5)/\operatorname{Hom}(L/(1-g$  $g(L^*, \mathbb{Z}_5) \cong 5^3$ . Combining these with Corollary 5.2, we have the following proposition.

**Proposition 6.16.** Stab<sub>Im  $\mu$ </sub>(V<sub>L</sub>(1))  $\cong$  5<sup>3</sup>. (2 × Ω<sub>3</sub>(5)).

Finally, we determine the structure of  $\operatorname{Im} \mu$ . Since  $\operatorname{Irr}(V_L^{\hat{g}}) \cong 5^5$  as abelian groups,  $\operatorname{Im} \mu$ is a subgroup of  $GO_5(5)$ . By [Wi09], we can compute the order of  $GO_5(5)$ . Combining this with Proposition 6.15, we see that  $|GO_5(5) : Im \mu| = 2$ . We summarize our results. The group Im  $\mu$  is a subgroup of GO<sub>5</sub>(5) satisfying the following:

- (1)  $|\text{GO}_5(5) : \text{Im } \mu| = 2;$
- (2) Im  $\mu$  acts transitively on  $S_{5C}$ ;
- (3) The stabilizer of an element in  $S_{5C}$  in Im  $\mu$  is isomorphic to  $5^3 \cdot (2 \times \Omega_3(5))$ .

There exist precisely three subgroups of  $GO_5(5)$  of index 2. These three groups are  $SO_5(5), 2 \times \Omega_5(5)$  and  $\Omega_5(5)$ . 2. By using MAGMA, we see that among the above three groups, there exists only one group which satisfies the condition (2) and (3). The group is  $2 \times \Omega_5(5)$ . Combining this with Proposition 6.13, we have the following theorem.

**Theorem 6.17.** Aut  $(V_{\Lambda_{5C}}^{\hat{g}}) \cong (5^2 . (5^2 : 2)) . (2 \times \Omega_5(5)).$ 

6.3. The case  $L = \Lambda_{11A}$ . The following lemma describes the structure of the quadratic space  $\mathcal{D}(L)$ .

**Lemma 6.18.** ([LS24, Lemma 7.5])  $(\mathcal{D}(L), q)$  is a non-singular 2-dimensional quadratic space over  $\mathbb{F}_{11}$  of (-)-type.

Let g be an automorphism of the Leech lattice which belongs to the conjugacy class 11A. By Theorem 3.5, we see that  $\operatorname{Irr}(V_L^{\hat{g}}) \cong 11^4$  as abelian groups. We need to determine which type the quadratic space  $(\operatorname{Irr}(V_L^{\hat{g}}), q)$  has. By Lemma 6.18, the quadratic space  $(\operatorname{Irr}(V_L^{\hat{g}}), q)$  has (-)-type. Moreover, by [Wi09, (3.27)], we can count the all singular vectors in  $(\operatorname{Irr}(V_L^{\hat{g}}), q)$ . Hence, we have the following proposition.

**Proposition 6.19.**  $(\operatorname{Irr}(V_L^{\hat{g}}), q)$  is of (-)-type. Hence, there are 1220 singular vectors in the quadratic space  $\operatorname{Irr}(V_L^{\hat{g}})$ .

Next, we consider the structure of Ker  $\mu$  in (5.1). By Theorem 5.1, we examine  $\operatorname{Hom}(L/(1-g)L^*,\mathbb{Z}_{11})$  and  $\{h \in C_{O(L)}(g) \mid h = 1 \text{ on } L^*/L\}/\langle g \rangle$ . By using MAGMA, we see that both groups are trivial. Hence we have the following proposition.

**Proposition 6.20.** Ker  $\mu$  is trivial. Hence the group homomorphism  $\mu$  is injective.

By Proposition 6.20, we see that  $\operatorname{Aut}(V_L^{\hat{g}}) \cong \operatorname{Im} \mu \subset \operatorname{GO}_4^-(11)$ . By using MAGMA, we see that  $\operatorname{Hom}(L/(1-g)L, \mathbb{Z}_{11}) \cong 11^2$  and  $C_{O(L)}(g)/\langle g \rangle \cong D_{12}$ . Hence we have the following proposition.

**Proposition 6.21.** Hom $(L/(1-g)L, \mathbb{Z}_{11}) \cong 11^2$  and  $C_{O(L)}(g)/\langle g \rangle \cong D_{12}$ . Hence we have Stab<sub>Aut  $(V_r^{\hat{g}})(V_L(1)) \cong 11^2$ .  $D_{12}$  and # Stab<sub>Aut  $(V_r^{\hat{g}})(V_L(1)) = 2^2 \cdot 3 \cdot 11^2$ .</sub></sub>

Next we consider the orbit of  $V_L(1)$ . By [LS24, Section 7], there exists  $\tau \in \operatorname{Aut}(V_L^{\hat{g}})$  such that

(6.1) 
$$V_L(1) \circ \tau$$
 is of twisted type.

Let  $X_{k,11A} = \{M \in \operatorname{Irr}(V_L^{\hat{g}}) \mid q(M) = 0 \text{ and } M \text{ is of } \hat{g}^k\text{-type}\}$ . Imitating [Wi09, (3.27), (3.28)], we can count the number of elements of  $X_{k,11A}$  for  $0 \leq k \leq 10$ . Hence we have the following lemma.

**Lemma 6.22.**  $\#X_{0,11A} = 11$  and  $\#X_{k,11A} = 121$  for  $1 \le k \le 10$ .

Combining Lemma 6.22 with (6.1) and Proposition 3.7, we have the following Lemma.

**Lemma 6.23.** Under the action of  $\operatorname{Im} \mu$ ,  $\#\operatorname{Orbit}(V_L(1)) \ge 122$ . Hence we have  $2 \le |\operatorname{GO}_4^-(11) : \operatorname{Im} \mu| \le 20$ .

Proof. By (6.1), we see that  $V_L(1) \circ \tau$  is of twisted type. Let it be of  $\hat{g}^s$ -type, where s is a positive integer such that  $1 \leq s \leq 10$ . Moreover, by Proposition 3.7, all irreducible  $V_L^{\hat{g}}$ modules of  $\hat{g}^s$ -type with same conformal weight are conjugate under the action of Aut  $(V_L^{\hat{g}})$ . By Lemma 6.22, since  $\#X_{s,11A} = 121$ , we have # Orbit $(V_L(1)) \geq 122$ . Hence, we see that  $\# \operatorname{Im} \mu \geq 122 \cdot \# \operatorname{Stab}_{\operatorname{Aut}(V_L^{\hat{g}})}(V_L(1))$ . By Proposition 6.21, we have  $\# \operatorname{Stab}_{\operatorname{Aut}(V_L^{\hat{g}})}(V_L(1)) =$  $2^2 \cdot 3 \cdot 11^2$ . By [Wi09, (3.32)], since  $\#\operatorname{GO}_4^-(11) = 2^5 \cdot 3 \cdot 5 \cdot 11^2 \cdot 61$ , we have  $|\operatorname{GO}_4^-(11) :$  $\operatorname{Im} \mu| \leq 20$ .

By Proposition 6.19, since  $\operatorname{Irr}(V_L^{\hat{g}})$  has 1220 singular vectors, we see that  $\# \operatorname{Im} \mu \leq 1220 \cdot \# \operatorname{Stab}_{\operatorname{Aut}(V_L^{\hat{g}})}(V_L(1)) = 1220 \cdot 2^2 \cdot 3 \cdot 11^2 = \# \operatorname{GO}_4^-(11)/2$ . This implies that  $2 \leq |\operatorname{GO}_4^-(11) : \operatorname{Im} \mu|$ .

Hence Im  $\mu$  is a subgroup of  $GO_4^-(11)$  satisfying the following:

- (1)  $2 \leq |\text{GO}_4^-(11) : \text{Im } \mu| \leq 20;$
- (2) The stabilizer of  $V_L(1)$  in Im  $\mu$  is isomorphic to  $11^2$ .  $D_{12}$ .

By using MAGMA, we can verify that there are precisely four subgroups of  $GO_4^-(11)$  satisfying the condition (1). These groups are  $\Omega_4^-(11), 2 \times \Omega_4^-(11), \Omega_4^-(11) \cdot 2$  and  $\Omega_4^-(11) \cdot 2$ . By using MAGMA, two groups of the these  $\Omega_4^-(11) \cdot 2$  have the same group structure. Moreover, by using MAGMA, we see that among the four groups, there exist two subgroups satisfying (2). The structure of the groups is  $\Omega_4^-(11) \cdot 2$ . Combining this with Proposition 6.20, we have the following theorem.

**Theorem 6.24.** Aut  $(V_{\Lambda_{11A}}^{\hat{g}}) \cong \Omega_4^-(11)$ . 2. Hence, Aut  $(V_{\Lambda_{11A}}^{\hat{g}})$  acts transitivity on all singular vectors in  $\operatorname{Irr}(V_{\Lambda_{11A}}^{\hat{g}})$ .

*Remark* 6.25. By [Wi09, (3.57)], since  $\Omega_4^-(11) \cong \text{PSL}_2(11^2)$ , we have  $\text{Aut}(V_{\Lambda_{11A}}^{\hat{g}}) \cong \text{PSL}_2(11^2)$ . 2

6.4. The case  $L = \Lambda_{23A}$ . Next we consider the case  $L = \Lambda_{23A}$ . By [LS24, Section 7], we see that  $\mathcal{D}(L) \cong 23$ . Let g be an automorphism of the Leech lattice which belongs to the conjugacy class 23A. By Theorem 3.5, we see that  $\operatorname{Irr}(V_L^{\hat{g}}) \cong 23^3$ .

Next, we consider the structure of Ker  $\mu$  in (5.1). By Theorem 5.1, we examine  $\operatorname{Hom}(L/(1-g)L^*, \mathbb{Z}_{23})$  and  $\{h \in C_{O(L)}(g) \mid h = 1 \text{ on } L^*/L\}/\langle g \rangle$ . By using MAGMA, we see that both groups are trivial. Hence we have the following proposition.

**Proposition 6.26.** Ker  $\mu$  is trivial. Hence the group homomorphism  $\mu$  is injective.

Hence we see that  $\operatorname{Aut}(V_L^{\hat{g}}) \cong \operatorname{Im} \mu \subset \operatorname{GO}_3(23)$ . By using MAGMA, we can verify that  $\operatorname{Hom}(L/(1-g)L, \mathbb{Z}_{23}) \cong 23$  and  $C_{O(L)}(g)/\langle g \rangle \cong 2$ . Hence we have the following proposition.

**Proposition 6.27.** Hom $(L/(1-g)L, \mathbb{Z}_{23}) \cong 23$  and  $C_{O(L)}(g)/\langle g \rangle \cong 2$ . Hence, we have  $\operatorname{Stab}_{\operatorname{Aut}(V_L^{\hat{g}})}(V_L(1)) \cong 23.2$  and  $\# \operatorname{Stab}_{\operatorname{Aut}(V_L^{\hat{g}})}(V_L(1)) = 2 \cdot 23.$ 

Next we consider the orbit of  $V_L(1)$ . By [LS24, Section 7], there exists  $\tau \in \operatorname{Aut}(V_L^{\hat{g}})$  such that

(6.2) 
$$V_L(1) \circ \tau$$
 is of twisted type.

Let  $X_{k,23A} = \{M \in \operatorname{Irr}(V_L^{\hat{g}}) \mid q(M) = 0 \text{ and } M \text{ is of } \hat{g}^k\text{-type}\}$ . Imitating [Wi09, (3.27), (3.28)], we can count the number of elements of  $X_{k,23A}$  for  $0 \leq k \leq 22$ . Hence we have the following lemma.

Lemma 6.28.  $\#X_{k,23A} = 23$  for  $0 \le k \le 22$ .

By imitating the discussion in the case 11A, we have the following lemma.

**Lemma 6.29.**  $\# \operatorname{Orbit}(V_L(1)) \ge 24$ . Hence, we have that  $|\operatorname{GO}_3(23) : \operatorname{Im} \mu| \le 22$ .

*Proof.* Imitating the proof of Lemma 6.23, we have  $\# \operatorname{Orbit}(V_L(1)) \ge 24$ . By Proposition 6.27, since  $\# \operatorname{Stab}_{\operatorname{Aut}(V_L^{\hat{g}})}(V_L(1)) = 2 \cdot 23$ , we see that  $\# \operatorname{Im} \mu \ge 24 \cdot \# \operatorname{Stab}_{\operatorname{Aut}(V_L^{\hat{g}})}(V_L(1)) = 2^4 \cdot 3 \cdot 23$ . By [Wi09, (3.30)], since  $\# \operatorname{GO}_3(23) = 2^5 \cdot 3 \cdot 11 \cdot 23$ , we have  $|\operatorname{GO}_3(23) : \operatorname{Im} \mu| \le \# \operatorname{GO}_3(23)/(2^4 \cdot 3 \cdot 23) = 22$ . □

Hence Aut  $(V_L^{\hat{g}})$  is a subgroup of GO<sub>3</sub>(23) satisfying the following:

- (1)  $|\text{GO}_3(23) : \text{Im } \mu| \le 22;$
- (2) The stabilizer of  $V_L(1)$  in Im  $\mu$  is isomorphic to 23.2.

By using MAGMA, we can verify that there are precisely five subgroups of  $GO_3(23)$  satisfying the condition (1). These groups are  $\Omega_3(23), 2 \times \Omega_3(23), SO_3(23), \Omega_3(23), 2$  and  $GO_3(23)$ . By using MAGMA, we see that among these five groups, there are two groups satisfying (2). The groups are  $\Omega_3(23), 2$  and  $GO_3(23)$ .

Remark 6.30. By using MAGMA, for  $\Omega_3(23)$ . 2 and  $GO_3(23)$ , we can count the number of the orbits on  $23^3$ . The number of the orbits of  $\Omega_3(23)$ . 2 (resp.  $GO_3(23)$ ) is 2 (resp. 1). For the group  $\Omega_3(23)$ . 2, the number of elements of each orbit is 264.

We determine which group Aut  $(V_L^{\hat{g}})$  is. To determine, we first examine the structure of the normalizer  $N_{\text{Aut}(V_L)}(\langle \hat{g} \rangle)/\langle \hat{g} \rangle$ . By using MAGMA, we see that  $N_{O(L)}(\langle g \rangle)/\langle g \rangle \cong 22$ . Moreover, by Proposition 6.27, we see  $\text{Hom}(L/(1-g)L, \mathbb{Z}_{23}) \cong 23$ . Combining these with Theorem 2.7 and Proposition 3.4, we have the following lemma.

Lemma 6.31. Stab<sub>Aut  $(V_L^{\hat{g}})$ </sub> ({ $V_L(i) \mid 0 \le i \le 22$ })  $\cong 23.22$ .

By using Lemma 6.31, we have the following lemma.

**Lemma 6.32.** Under the action of the stabilizer  $\operatorname{Stab}_{\operatorname{Aut}(V_L^{\hat{g}})}(\{V_L(i) \mid 0 \le i \le 22\})$  on  $\{V_L(i) \mid 0 \le i \le 22\}$ , we have  $\#\operatorname{Orbit}(V_L(1)) = 11$ .

*Proof.* By Lemma 6.31, we have  $\# \operatorname{Stab}_{\operatorname{Aut}(V_L^{\hat{g}})}(\{V_L(i) \mid 0 \le i \le 22\}) = 506$ . Since  $\operatorname{Stab}_{\operatorname{Aut}(V_L^{\hat{g}})}(V_L(1)) \subset \operatorname{Stab}_{\operatorname{Aut}(V_L^{\hat{g}})}(\{V_L(i) \mid 0 \le i \le 22\})$ , we have  $\# \operatorname{Orbit}(V_L(1)) = 506/46 = 11$ , as desired. □

Finally we have the following theorem.

**Theorem 6.33.** Aut  $(V_{\Lambda_{23A}}^{\hat{g}}) \cong \Omega_3(23)$ . 2. Hence, by Remark 6.30, under the action of Aut  $(V_{\Lambda_{23A}}^{\hat{g}})$ , the set of all singular vectors in  $\operatorname{Irr}(V_{\Lambda_{23A}}^{\hat{g}})$  is divided into two orbits which have same number of elements.

Proof. For contradiction, suppose that  $\operatorname{Aut}(V_L^{\hat{g}}) = \operatorname{GO}_3(23)$ . The center of  $\operatorname{GO}_3(23)$ is  $\{1, -1\}$ , where -1 is the diagonal matrix whose all diagonal entries are -1. Since  $\{V_L(i) \mid 0 \leq i \leq 22\}$  is a subspace of  $\operatorname{Irr}(V_{\Lambda_{23A}}^{\hat{g}})$ , the map -1 preserves  $\{V_L(i) \mid 0 \leq i \leq 22\}$ . Hence we see that -1 is in  $\operatorname{Stab}_{\operatorname{Aut}(V_L^{\hat{g}})}(\{V_L(i) \mid 0 \leq i \leq 22\})$ .

On the other hand, by Lemma 6.32, under the action of  $\operatorname{Stab}_{\operatorname{Aut}(V_L^{\hat{g}})}(\{V_L(i) \mid 0 \leq i \leq 22\})$ , we have  $\# \operatorname{Orbit}(V_L(1)) = 11$ . Since the stabilizer  $\operatorname{Stab}_{\operatorname{Aut}(V_L^{\hat{g}})}(\{V_L(i) \mid 0 \leq i \leq 22\})$  has the element -1 and -1 is fixed-point free, its orbits which do not contain the element  $V_L(0)$  have even number of elements. This is contradiction, and we have the desired result.

*Remark* 6.34. By [Wi09, (3.57)], since  $\Omega_3(23) \cong \text{PSL}_2(23)$ , we have Aut  $(V_{\Lambda_{23A}}^{\hat{g}}) \cong \text{PSL}_2(23)$ . 2

#### 7. Conclusion

Throughout Section 6, we determined the automorphism groups of the orbifold VOAs. In conclusion, we have the following theorem.

**Theorem 7.1.** Let  $(g, L) \in \{(3C, \Lambda_{3C}), (5C, \Lambda_{5C}), (11A, \Lambda_{11A}), (23A, \Lambda_{23A})\}$  and let  $\hat{g}$  be a lift of g in Subsection 2.4. Then the automorphism groups of orbifold VOAs  $V_L^{\hat{g}}$  are following:

- Aut  $(V_{\Lambda_{3C}}^{\hat{g}}) \cong (3^4 \cdot (3^4 : 2)) \cdot (\Omega_7(3) \cdot 2));$
- Aut  $(V_{\Lambda_{5C}}^{\hat{g}}) \cong (5^2 . (5^2 : 2)) . (2 \times \Omega_5(5));$
- Aut  $(V_{\Lambda_{11A}}^{\hat{g}}) \cong \Omega_4^-(11)$ . 2;
- Aut  $(V_{\Lambda_{234}}^{\hat{g}}) \cong \Omega_3(23)$ . 2.

Acknowledgment. I would like to express my gratitude to my supervisor Professor Koichi Betsumiya. I thank Professor Ching Hung Lam for encouragement to consider the problem in this paper and for helpful advice. I also thank Professor Hiroki Shimakura for useful comments about this problem and for helpful advice.

#### References

- [ALY18] T. Abe, C.H. Lam and H. Yamada, Abe, Extensions of tensor products of  $\mathbb{Z}_p$ -orbifold models of the lattice vertex operator algebra  $V_{\sqrt{2}A_{p-1}}$ , J. Algebra **510** (2018), 24–51.
- [ATLAS] J.H. Conway, R.T. Curtis, S.P. Norton, R.A. Parker and R.A. Wilson, ATLAS of finite groups. Clarendon Press, Oxford, 1985.
- [BCP97] W. Bosma, J. Cannon and C. Playoust, The Magma algebra system I: The user language, J. Symbolic Comput. 24 (1997), 235–265.
- [BLS24] K. Betsumiya, C.H. Lam and H. Shimakura, Automorphism groups of cyclic orbifold vertex operator algebras associated with the Leech lattice and some non-prime isometries, *Isr.J. Math.* 259 (2024), no.2, 621-650.
- [CM] S. Carnahan and M. Miyamoto, Regularity of fixed-point vertex operator subalgebras; arXiv:1603.05645.
- [CLS18] H.Y. Chen, C.H. Lam and H. Shimakura, On Z<sub>3</sub>-orbifold construction of the Moonshine vertex operator algebra, Math. Z. 288 (2018), no. 1-2, 75-100.
- [DLM00] C. Dong, H. Li, and G. Mason, Modular-invariance of trace functions in orbifold theory and generalized Moonshine, *Comm. Math. Phys.* 214 (2000), 1–56.
- [DN99] C. Dong and K. Nagatomo, Automorphism groups and twisted modules for lattice vertex operator algebras, in Recent developments in quantum affine algebras and related topics (Raleigh, NC, 1998), 117–133, Contemp. Math., 248, Amer. Math. Soc., Providence, RI, 1999.
- [DRX17] C. Dong, L. Ren and F. Xu, On Orbifold Theory, Adv. Math. 321 (2017), 1–30.
- [EMS20] J. van Ekeren, S. Möller and N. Scheithauer, Construction and classification of holomorphic vertex operator algebras, J. Reine Angew. Math., 759 (2020), 61–99.
- [FHL93] I.B. Frenkel, Y. Huang and J. Lepowsky, On axiomatic approaches to vertex operator algebras and modules, *Mem. Amer. Math. Soc.* **104** (1993), viii+64 pp.
- [FLM88] I. Frenkel, J. Lepowsky and A. Meurman, Vertex operator algebras and the Monster, Pure and Appl. Math., Vol.134, Academic Press, Boston, 1988.
- [Gr02] L.C. Grove, Classical Groups and Geometric Algebra. Graduate Studies in Mathematics, 39. American Mathematical Society, Providence, RI, 2002.
- [HL95] Y.Z. Huang and J. Lepowsky, A theory of tensor product for module category of a vertex operator algebra, III, J. Pure Appl. Algebra, 100 (1995), 141–171.
- [La20] C.H. Lam, Cyclic orbifolds of lattice vertex operator algebras having group-like fusions, Lett. Math. Phys. 110 (2020), 1081–1112.
- [La22+] C.H. Lam, Some observations about the automorphism groups of certain orbifold vertex operator algebras, to appear in RIMS Kôkyûroku Bessatsu.
- [LS24] C.H. Lam and H. Shimakura, Extra automorphisms of cyclic orbifolds of lattice vertex operator algebras, Jornal of pure and Applied Algebra 228 (2024), Article no. 107454.

- [LS20] C.H. Lam and H. Shimakura, On orbifold constructions associated with the Leech lattice vertex operator algebra, Math. Proc. Cambridge Philos. Soc. 168 (2020), 261–285.
- [LY14] C.H. Lam and H. Yamauchi, On 3-transposition groups generated by  $\sigma$ -involutions associated to c = 4/5 Virasoro vectors, J. Algebra, 416 (2014), 84-121.
- [Mi15] M. Miyamoto, C<sub>2</sub>-cofiniteness of cyclic-orbifold models, Comm. Math. Phys. 335 (2015), 1279– 1286.
- [Sh04] H. Shimakura, The automorphism group of the vertex operator algebra  $V_L^+$  for an even lattice L without roots, J. Algebra **280** (2004), 29–57.
- [Wi09] R.A. Wilson, The finite simple groups, Graduate Texts in Mathematics, 251, Springer, London, 2009.

(T. Kondo) Graduate School of Science and Technology, Hirosaki University, Hirosaki 036-8561, Japan

Email address: h23ms107@hirosaki-u.ac.jp